

MORI DREAM SPACES AS FINE MODULI OF QUIVER REPRESENTATIONS

ALASTAIR CRAW AND DOROTHY WINN

ABSTRACT. We construct Mori Dream Spaces as fine moduli spaces of representations of bound quivers, thereby extending results of Craw–Smith [6] beyond the toric case. Any collection of effective line bundles $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ on a Mori Dream Space X defines a bound quiver of sections and a map from X to a toric quiver variety $|\mathcal{L}|$ called the multigraded linear series. We provide necessary and sufficient conditions for this map to be a closed immersion and, under additional assumptions on \mathcal{L} , the image realises X as the fine moduli space of ϑ -stable representations of the bound quiver. As an application, we show how to reconstruct del Pezzo surfaces from a full, strongly exceptional collection of line bundles.

CONTENTS

1. Introduction	1
2. Quivers of sections on Mori Dream Spaces	4
3. Morphism to the multigraded linear series	8
4. Fine moduli of bound quiver representations	13
5. Reconstructing del Pezzo surfaces from a tilting bundle	21
References	25

1. INTRODUCTION

Mori Dream Spaces and their Cox rings have been the subject of a great deal of interest since their introduction by Hu–Keel [11] over a decade ago. From the geometric side, these varieties enjoy the property that all operations of the Mori programme can be carried out by variation of GIT quotient, while from the algebraic side, obtaining an explicit presentation of the Cox ring is an interesting problem in itself. Examples of Mori Dream Spaces include \mathbb{Q} -factorial projective toric varieties, spherical varieties and log Fano varieties of arbitrary dimension. In this paper we use the representation theory of quivers to study multigraded linear series on Mori Dream Spaces. Our main results construct Mori Dream Spaces as fine moduli spaces of ϑ -stable representations of bound quivers for a special stability parameter ϑ , thereby extending results of Craw–Smith [6] for projective toric varieties.

Let X be a projective variety and $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ a collection of distinct, effective line bundles on X . The associated *multigraded linear series* $|\mathcal{L}|$ is a smooth projective toric variety that provides a multigraded analogue of the classical linear series of a single line bundle. To construct $|\mathcal{L}|$ one first defines the bound quiver of sections of \mathcal{L} to be a finite, acyclic

MSC 2010: Primary 14D22, 16G20; Secondary 14F05, 16E35, 16S38.

quiver Q together with an ideal of relations $J_{\mathcal{L}}$ in the path algebra $\mathbb{k}Q$ for which $\mathbb{k}Q/J_{\mathcal{L}}$ is isomorphic to $\text{End}(\bigoplus_{0 \leq i \leq r} L_i)$. Setting aside the ideal of relations for now, the multigraded linear series $|\mathcal{L}|$ is defined to be the toric quiver variety obtained as the fine moduli space of ϑ -stable representations of Q with dimension vector $(1, \dots, 1)$ for the special stability parameter $\vartheta = (-r, 1, \dots, 1)$. Paths in the quiver arise from spaces of sections of the form $H^0(X, L_j \otimes L_i^{-1})$ for $0 \leq i, j \leq r$, and evaluating these sections defines a morphism

$$(1.1) \quad \varphi_{|\mathcal{L}|}: X \longrightarrow |\mathcal{L}|$$

when each L_i is basepoint-free. In the case where X is a projective toric variety, Craw–Smith [6] constructed the quiver of sections using only torus-invariant sections of the bundles $L_j \otimes L_i^{-1}$, and described precisely the image of $\varphi_{|\mathcal{L}|}$. The bound quiver of sections and a morphism $\varphi_{|\mathcal{L}|}$ are constructed in general by Craw [5, Theorem 5.4], but one cannot study the image without choosing bases for the appropriate spaces of sections.

Our first main result solves this problem for Mori Dream Spaces by choosing a suitable spanning set for each space of sections. Every Mori Dream Space X admits a closed immersion into a \mathbb{Q} -factorial projective toric variety \tilde{X} , such that each line bundle on X is the restriction of a reflexive sheaf of rank one on \tilde{X} . In particular, the collection \mathcal{L} on X can be lifted uniquely to a collection $\tilde{\mathcal{L}} = (\mathcal{O}_{\tilde{X}}, E_1, \dots, E_r)$ of rank one reflexive sheaves on \tilde{X} . While the spaces $H^0(X, L_j \otimes L_i^{-1})$ have no preferred basis, the restriction to X of the torus-invariant sections of the sheaf $(E_j \otimes E_i^{\vee})^{\vee\vee}$ provide a spanning set for $H^0(X, L_j \otimes L_i^{-1})$ that we use to build the paths in the quiver Q from vertex i to vertex j . That is, we *define* the quiver Q using differences of sections of bundles from the collection $\tilde{\mathcal{L}}$, and we introduce an ideal of relations $J_{\mathcal{L}}$ in $\mathbb{k}Q$ for which $\mathbb{k}Q/J_{\mathcal{L}} \cong \text{End}(\bigoplus_{0 \leq i \leq r} L_i)$. Our first main result describes the rational map

$$\varphi_{|\mathcal{L}|}: X \dashrightarrow |\mathcal{L}|$$

obtained by evaluating sections, and establishes the link between the collection $(\mathcal{W}_0, \dots, \mathcal{W}_r)$ of tautological line bundles on the fine moduli space $|\mathcal{L}|$ and the collection \mathcal{L} of line bundles on X (see Section 3.2 for the proof):

Theorem 1.1. *For a collection $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ of distinct, effective line bundles on X , the map $\varphi_{|\mathcal{L}|}: X \dashrightarrow |\mathcal{L}|$ is a morphism if and only if each L_i is basepoint-free. Moreover, the image is presented explicitly as a geometric quotient, and the tautological bundles on $|\mathcal{L}|$ satisfy $\varphi_{|\mathcal{L}|}^*(\mathcal{W}_i) = L_i$.*

In addition, we provide a necessary and sufficient condition (that is easy to implement) for the morphism $\varphi_{|\mathcal{L}|}: X \rightarrow |\mathcal{L}|$ to be a closed immersion, thereby improving upon a sufficient condition in the toric case by Craw–Smith [6, Corollary 4.10]. A straightforward application of multigraded regularity introduced by Hering–Schenck–Smith [9] (see also Maclagan–Smith [14]) provides an efficient way to exhibit many collections that give rise to closed immersions. If each L_i on X is the restriction of a basepoint-free line bundle on \tilde{X} then the morphism from Theorem 1.1 is simply the restriction of that from [6, Theorem 1]. This is rarely the case however, because the nef cone of X is typically the union of the nef cones of a finite collection of ambient toric varieties.

Our second main result is more algebraic, and provides a fine moduli description of X via the noncommutative algebra $A := \mathbb{k}Q/J_{\mathcal{L}}$. Since $\mathbb{k}Q$ is constructed from sections of sheaves on

the ambient toric variety rather than sections of sheaves on X , one does not expect a strong link between X and the multigraded linear series until the geometry is influenced by the ideal of relations $J_{\mathcal{L}}$. To this end, the ideal $J_{\mathcal{L}}$ defines an ideal $I_{\mathcal{L}}$ in the Cox ring of $|\mathcal{L}|$, and the subscheme cut out by this ideal is the fine moduli space $\mathcal{M}_{\vartheta}(\text{mod-}A)$ of ϑ -stable A -modules with dimension vector $(1, \dots, 1)$. This subscheme contains the image of the morphism $\varphi_{|\mathcal{L}|}$ from Theorem 1.1. This inclusion is proper in general, but by saturating $I_{\mathcal{L}}$ with the irrelevant ideal for the GIT quotient construction of $|\mathcal{L}|$ and by comparing the result with the ideal I_Q that cuts out the image of $\varphi_{|\mathcal{L}|}$, we obtain the following (see Section 4.4):

Theorem 1.2. *For any Mori Dream Space X , there exist (many) collections \mathcal{L} on X such that the morphism $\varphi_{|\mathcal{L}|}: X \rightarrow |\mathcal{L}|$ identifies X with the fine moduli space $\mathcal{M}_{\vartheta}(\text{mod-}A)$, and the tautological line bundles on $\mathcal{M}_{\vartheta}(\text{mod-}A)$ coincide with the line bundles of \mathcal{L} .*

The analogous result for toric varieties [6, Theorem 1.2] relies heavily on lattice-theoretic and combinatorial techniques that are unavailable for Mori Dream Spaces. Our proof of Theorem 1.2 combines a careful analysis of paths in the quiver of sections together with several invocations of multigraded regularity.

More generally, whenever the morphism $\varphi_{|\mathcal{L}|}: X \rightarrow |\mathcal{L}|$ is a closed immersion it identifies X with $\mathcal{M}_{\vartheta}(\text{mod-}A)$ when the saturation of $I_{\mathcal{L}}$ by the irrelevant ideal coincides with the ideal I_Q . These ideals can be computed explicitly in any given example, so it is possible to check directly whether Theorem 1.2 holds (subject to computational limitations). The resulting geometric quotient constructions of X are new, and while it is impossible to improve upon the Hu–Keel construction of X from the birational point of view, it is sometimes possible to encode more refined information of X via \mathcal{L} , such as its bounded derived category of coherent sheaves

As an application of our results we illustrate a quiver moduli construction that encodes the bounded derived category of coherent sheaves for a pair of del Pezzo surfaces in which \mathcal{L} is chosen to be a full, strongly exceptional collection of basepoint-free line bundles, that is, the line bundles in \mathcal{L} freely generate the bounded derived category of coherent sheaves on X . In each case, we deduce from Theorem 1.1 and its corollaries that $\varphi_{|\mathcal{L}|}: X \rightarrow |\mathcal{L}|$ is a closed immersion and, moreover, we compute that the saturation of $I_{\mathcal{L}}$ by the irrelevant ideal coincides with the ideal I_Q as in Theorem 1.2. It follows in each case that the del Pezzo surface X is isomorphic to the fine moduli space $\mathcal{M}_{\vartheta}(\text{mod-}A)$, and the tautological bundles on the moduli space correspond under this isomorphism to the line bundles in \mathcal{L} . Until now it was known only that each del Pezzo surface X is isomorphic to a connected component of a fine moduli space $\mathcal{M}_{\vartheta}(\text{mod-}A)$, see Bergman–Proudfoot [3].

We now describe the structure of the paper. Section 2 defines the bound quiver of sections for collections of line bundles on a Mori Dream Space. The morphism to the multigraded linear series and the geometric results leading to Theorem 1.1 are introduced in Section 3, while the ideals of relations and the algebraic results leading to Theorem 1.2 appear in Section 4. Finally, our application to tilting bundles on del Pezzo surfaces is presented in Section 5.

Conventions. Write \mathbb{k} for an algebraically closed field of characteristic zero, \mathbb{k}^{\times} for the one-dimensional algebraic torus over \mathbb{k} , and \mathbb{N} for the semigroup of nonnegative integers. Given a vector space V , we write $\mathbb{P}^*(V)$ for the projective space of hyperplanes in V . For $d > 0$ and $\mathbf{u} := (u_1, \dots, u_d) \in \mathbb{N}^d$ we adopt multiindex notation $x^{\mathbf{u}} := \prod_{1 \leq i \leq d} x_i^{u_i} \in \mathbb{k}[x_1, \dots, x_d]$.

Acknowledgements. Special thanks to Tarig Abdelgadir and Arend Bayer for useful conversations, and to the anonymous referee for helpful comments. Thanks also to Jürgen Hausen, Markus Perling and Diane Maclagan. This work forms part of the second author's EPSRC-funded PhD thesis. The first author was supported in part by EPSRC grant EP/G004048.

2. QUIVERS OF SECTIONS ON MORI DREAM SPACES

We begin by establishing our notation and introducing the bound quiver of sections for a collection of line bundles on a Mori Dream Space. These bound quivers encode the endomorphism algebra of the direct sum of the sheaves in the collection.

2.1. Mori Dream Spaces. Let X be a projective \mathbb{Q} -factorial variety with divisor class group $\text{Cl}(X)$ that is finitely generated and free of rank ρ . Let D_1, \dots, D_ρ be Weil divisors whose classes provide an integral basis of $\text{Cl}(X)$. Assume that the \mathbb{Z}^ρ -graded *Cox ring*

$$\text{Cox}(X) := \bigoplus_{(m_1, \dots, m_\rho) \in \mathbb{Z}^\rho} H^0(X, \mathcal{O}_X(m_1 D_1 + \dots + m_\rho D_\rho))$$

is a finitely generated \mathbb{k} -algebra, where the multiplication is induced by multiplication of global sections. A different choice of integral basis for the divisor class group yields (non-canonically) an isomorphic Cox ring for X . Since $\text{Cl}(X)$ is a finitely generated and free abelian group, $\text{Cox}(X)$ is a unique factorisation domain by Elizondo–Kurano–Watanabe [7]. The main result of Hu–Keel [11, Proposition 2.9] shows that finite generation of $\text{Cox}(X)$ is equivalent to X being a *Mori Dream Space* (see [11] for the definition that justifies the terminology).

Choose once and for all a presentation

$$(2.1) \quad \text{Cox}(X) = \mathbb{k}[x_1, \dots, x_d]/I_X$$

for some $d \in \mathbb{N}$. The \mathbb{Z}^ρ -grading from $\text{Cox}(X)$ can be lifted via the quotient map

$$(2.2) \quad \tau: \mathbb{k}[x_1, \dots, x_d] \longrightarrow \text{Cox}(X).$$

to obtain a \mathbb{Z}^ρ -grading of $\mathbb{k}[x_1, \dots, x_d]$ that makes τ a \mathbb{Z}^ρ -graded homomorphism of \mathbb{k} -algebras. This defines a semigroup homomorphism $\deg: \mathbb{N}^d \rightarrow \mathbb{Z}^\rho$ and an action of the algebraic torus $T := \text{Hom}(\mathbb{Z}^\rho, \mathbb{k}^\times)$ on $\mathbb{A}_{\mathbb{k}}^d$. Choose $\chi \in T^* = \mathbb{Z}^\rho$ such that every χ -semistable point of $\mathbb{A}_{\mathbb{k}}^d$ is χ -stable and, moreover, such that a χ -stable point exists (that is, we assume χ lies in an open GIT chamber for the T -action on $\mathbb{A}_{\mathbb{k}}^d$). Then the radical monomial ideal

$$B = \text{rad}(x^{\mathbf{u}} \in \mathbb{k}[x_1, \dots, x_d] : \deg(\mathbf{u}) = m\chi \text{ for some } m > 0)$$

cuts out the χ -unstable locus in $\mathbb{A}_{\mathbb{k}}^d$, and the geometric quotient $\mathbb{A}_{\mathbb{k}}^d //_{\chi} T := (\mathbb{A}_{\mathbb{k}}^d \setminus \mathbb{V}(B))/T$ of the open subscheme of χ -stable points in $\mathbb{A}_{\mathbb{k}}^d$ by T is a projective \mathbb{Q} -factorial toric variety.

The action of T restricts to an action on $\text{Spec}(\text{Cox}(X)) \cong \mathbb{V}(I_X)$ since τ is \mathbb{Z}^ρ -graded. Choose $\chi \in T^*$ that is ample on X (so χ -stable points exist), and assume that every χ -semistable point of $\mathbb{A}_{\mathbb{k}}^d$ is χ -stable. The χ -unstable locus in $\mathbb{V}(I_X)$ is cut out by the ideal $B_X := \text{rad}(I_X + B)$. Since χ is ample on X , Hu–Keel [11, Proposition 2.11] implies that the geometric quotient $\mathbb{V}(I_X) //_{\chi} T := (\mathbb{V}(I_X) \setminus \mathbb{V}(B_X))/T$ of the open subscheme of χ -stable points by the action of T is isomorphic to X . Since $\mathbb{V}(I_X) \subseteq \mathbb{A}_{\mathbb{k}}^d$ is closed, there is a closed immersion

$$\mathbb{V}(I_X) //_{\chi} T \hookrightarrow \mathbb{A}_{\mathbb{k}}^d //_{\chi} T$$

that embeds the Mori Dream Space X into what we call the *ambient toric variety* $\tilde{X}_\chi := \mathbb{A}_{\mathbb{k}}^d //_\chi T$. In fact a choice is required here: the GIT chamber decomposition for the action of T on $\mathbb{A}_{\mathbb{k}}^d$ refines that for the action on $\text{Spec}(\text{Cox}(X))$, so the nef cone of X is the union of nef cones of finitely many ambient toric varieties. Nevertheless, any two such are isomorphic in codimension-one, and our construction requires only knowledge of the divisors in \tilde{X}_χ . Hereafter we choose one such ambient toric variety $\tilde{X} := \tilde{X}_\chi$ and suppress the dependence on χ .

Every Mori Dream Space can be constructed from a bunched ring in the sense of Berchtold–Hausen [2, Definition 2.10, Theorem 4.2]. It follows that the restriction of rank one reflexive sheaves from the ambient toric variety \tilde{X} to X induces an isomorphism

$$(2.3) \quad \psi: \text{Cl}(\tilde{X}) \longrightarrow \text{Cl}(X) = \mathbb{Z}^p$$

on divisor class groups, see [2, Proposition 5.2].

Example 2.1. Let X_4 be a del Pezzo surface obtained as the blow-up of $\mathbb{P}_{\mathbb{k}}^2$ at four points in general position. The Picard group \mathbb{Z}^5 has a basis given by H , the pullback to X_4 of the hyperplane class on $\mathbb{P}_{\mathbb{k}}^2$, together with the four exceptional curves R_1, R_2, R_3, R_4 . The semigroup homomorphism $\deg: \mathbb{N}^{10} \rightarrow \mathbb{Z}^5$ obtained as multiplication by the matrix

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & -1 & -1 & -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -1 & 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 & -1 & 0 & -1 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 & 0 & -1 & -1 \end{bmatrix}$$

induces the \mathbb{Z}^5 -grading of $\mathbb{k}[x_1, \dots, x_{10}]$, and the T -homogeneous ideal

$$I_{X_4} := \begin{pmatrix} x_2x_5 - x_3x_6 + x_4x_7, & x_1x_5 - x_3x_8 + x_4x_9 \\ x_1x_6 - x_2x_8 + x_4x_{10}, & x_1x_7 - x_2x_9 + x_3x_{10}, & x_5x_{10} - x_6x_9 + x_7x_8 \end{pmatrix}$$

determines $\text{Cox}(X_4) = \mathbb{k}[x_1, \dots, x_{10}]/I_{X_4}$ following Batyrev–Popov [1]. The ample linearisation $-K_{X_4} = 3H - R_1 - R_2 - R_3 - R_4$ is a natural choice to construct an ambient toric variety, but as Laface–Velasco [13, Example 2.11] note, this defines a non- \mathbb{Q} -factorial toric quotient so it must lie in a GIT wall for the action of T on $\mathbb{A}_{\mathbb{k}}^{10}$. However, the same authors also note that $\chi = 11H - 5R_1 - 3R_2 - 2R_3 - R_4 \in \mathbb{Z}^5$ is ample on X and $\mathbb{A}_{\mathbb{k}}^{10} //_\chi T$ is \mathbb{Q} -factorial so it lies in an open GIT chamber for the action of T on $\mathbb{A}_{\mathbb{k}}^{10}$. We therefore choose $\tilde{X}_4 := \mathbb{A}_{\mathbb{k}}^{10} //_\chi T$.

2.2. Quivers and path algebras. We now introduce notation for quivers and path algebras.

Let Q be a finite connected quiver with vertex set Q_0 , arrow set Q_1 , and maps $h, t: Q_1 \rightarrow Q_0$ indicating the head and tail of each arrow. The characteristic functions $\chi_i: Q_0 \rightarrow \mathbb{Z}$ for $i \in Q_0$ and $\chi_a: Q_1 \rightarrow \mathbb{Z}$ for $a \in Q_1$ form the standard integral bases of the vertex space \mathbb{Z}^{Q_0} and the arrow space \mathbb{Z}^{Q_1} respectively. A nontrivial path in Q is a sequence of arrows $p = a_k \cdots a_1$ with $h(a_j) = t(a_{j+1})$ for $1 \leq j < k$. Set $t(p) = t(a_1)$, $h(p) = h(a_k)$ and $\text{supp}(p) = \{a_1, \dots, a_k\}$. Each $i \in Q_0$ gives a trivial path e_i where $t(e_i) = h(e_i) = i$. The path algebra $\mathbb{k}Q$ is the \mathbb{k} -algebra whose underlying \mathbb{k} -vector space has a basis of paths in Q , where the product of basis elements is the basis element defined by concatenation of the paths if possible, or zero otherwise. A cycle is a path p with $t(p) = h(p)$, and a quiver is acyclic if every cycle is a trivial path e_i for some vertex $i \in Q_0$. A vertex $i \in Q_0$ is a source of Q if it is not the head of an arrow. A spanning tree

in Q is a connected acyclic subquiver whose vertex set coincides with that of Q , and a spanning tree has root at $i \in Q_0$ if i is the unique source of the tree.

A relation in a quiver Q is a \mathbb{k} -linear combination of paths that share the same head and tail. Thus, each relation is of the form $\sum_{p \in \Gamma} c_p p$ for $c_p \in \mathbb{k}$, where Γ is a finite set of paths that share the same head and the same tail. We do not assume that paths in Γ traverse at least two arrows. Any set of relations generates a two-sided ideal $J \subseteq \mathbb{k}Q$, and the pair (Q, J) is called a quiver with relations or, more simply, a *bound quiver*.

2.3. Quivers of sections. For $r \geq 0$, consider a collection of distinct line bundles

$$\mathcal{L} := (L_0, L_1, \dots, L_r) \subset \mathbb{Z}^\rho = \text{Cl}(X)$$

on the Mori Dream Space X , where $L_0 = \mathcal{O}_X$ and L_1, \dots, L_r are effective. For $0 \leq i \leq r$, define $E_i \in \psi^{-1}(L_i)$ using the isomorphism $\psi: \text{Cl}(\tilde{X}) \rightarrow \text{Cl}(X) = \mathbb{Z}^\rho$ from (2.3) to obtain a collection

$$\tilde{\mathcal{L}} := (E_0, E_1, \dots, E_r)$$

of distinct rank one reflexive sheaves on an ambient toric variety \tilde{X} , where $E_0 = \mathcal{O}_{\tilde{X}}$. Since \tilde{X} is normal, each sheaf E_i is of the form $\mathcal{O}_{\tilde{X}}(D)$ for some $D \in \text{Cl}(\tilde{X})$. For $0 \leq i \leq r$, define $D'_i \in \text{Cl}(\tilde{X})$ such that $E_i = \mathcal{O}_{\tilde{X}}(D'_i)$. Writing $E^\vee := \mathcal{H}om_{\mathcal{O}_{\tilde{X}}}(E, \mathcal{O}_{\tilde{X}})$ for the dual sheaf gives

$$(2.4) \quad (E_j \otimes E_i^\vee)^{\vee\vee} \cong \mathcal{O}_{\tilde{X}}(D'_j - D'_i)$$

for $0 \leq i, j \leq r$, see for example Cox–Little–Schenck [4, Proposition 8.0.6] (compare also Example 8.0.5 from *loc.cit.* to see the necessity for using the double dual). For $0 \leq i, j \leq r$, we say that a torus-invariant section $s \in H^0(\tilde{X}, (E_j \otimes E_i^\vee)^{\vee\vee})$ is *irreducible* if the section does not lie in the image of the multiplication map

$$(2.5) \quad H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(D'_j - D'_k)) \otimes_{\mathbb{k}} H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(D'_k - D'_i)) \longrightarrow H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(D'_j - D'_i))$$

for any $k \neq i, j$, where we use isomorphism (2.4). The following extends the notion of a quiver of sections for a collection of line bundles on a projective toric variety due to Craw–Smith [6].

Definition 2.2. The *quiver of sections* of the collection \mathcal{L} on X is defined to be the quiver Q with vertex set $Q_0 = \{0, \dots, r\}$, and where the arrows from i to j correspond to the irreducible sections in $H^0(\tilde{X}, (E_j \otimes E_i^\vee)^{\vee\vee})$.

Remark 2.3. (1) For $0 \leq i, j \leq r$, the space of sections of the sheaf from (2.4) is a subspace of the ring $\text{Cox}(\tilde{X}) = \mathbb{k}[x_1, \dots, x_d]$, and the restriction of the map τ from (2.2) to this subspace is a surjective \mathbb{k} -linear map

$$\tau_{ij}: H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(D'_j - D'_i)) \longrightarrow H^0(X, L_j \otimes L_i^{-1}) \cong \text{Hom}_{\mathcal{O}_X}(L_i, L_j).$$

Thus, the image under $\tau_{i,j}$ of the torus-invariant sections of $\mathcal{O}_{\tilde{X}}(D'_j - D'_i)$ provide a spanning set for the space $\text{Hom}_{\mathcal{O}_X}(L_i, L_j)$ of primary interest, see Proposition 2.6 below. We abuse terminology by calling Q the ‘quiver of sections of \mathcal{L} ’ even though paths in Q from i to j are not constructed from a basis of $\text{Hom}(L_i, L_j)$ as in the literature [5, 6].

(2) In Definition 2.2 we write the sheaf from (2.4) as $(E_j \otimes E_i^\vee)^{\vee\vee}$ to express it directly in terms of the collection $\tilde{\mathcal{L}}$. Of course, if each E_i is invertible then we can write

$$(2.6) \quad H^0(\tilde{X}, (E_j \otimes E_i^\vee)^{\vee\vee}) \cong \text{Hom}_{\mathcal{O}_{\tilde{X}}}(E_i, E_j)$$

and Q is the quiver of sections of the collection $\widetilde{\mathcal{L}}$ on \widetilde{X} as defined in [6].

- (3) Composing arrows in Q translates into multiplication of torus-invariant sections. That is, an arrow from i to k defined by a section of $(E_k \otimes E_i^\vee)^{\vee\vee}$ composed with an arrow from k to j defined by a section of $(E_j \otimes E_k^\vee)^{\vee\vee}$ gives the path from i to j defined by the product of these sections under the multiplication map (2.5).
- (4) Definition 2.2 depends a priori on the choice of ambient toric variety \widetilde{X} . However, given a presentation of the Cox ring as in (2.1), any two such ambient toric varieties are isomorphic in codimension-one, so Q is independent of the choice of \widetilde{X} .

Lemma 2.4. *The quiver of sections Q is connected, acyclic, and $0 \in Q_0$ is the unique source.*

Proof. We first suppose that Q admits a nontrivial cycle, so there exists $i \neq j$ such that both $H^0((E_j \otimes E_i^\vee)^{\vee\vee})$ and $H^0((E_i \otimes E_j^\vee)^{\vee\vee})$ are nonzero. Remark 2.3(3) implies that the product of these nonzero sections gives a nonconstant section of $\mathcal{O}_{\widetilde{X}}$, but \widetilde{X} is projective, so Q is acyclic after all. For $i \in Q_0$, the space $H^0((E_i \otimes E_0^\vee)^{\vee\vee}) \cong H^0(E_i)$ has a torus-invariant element since E_1, \dots, E_r effective and $E_0 \cong \mathcal{O}_{\widetilde{X}}$, giving rise to a path in Q from 0 to $i \in Q_0$ as required. \square

The quiver of sections depends purely on the collection of reflexive sheaves $\widetilde{\mathcal{L}}$ on \widetilde{X} , but we aim to encode information about the collection of line bundles \mathcal{L} on the Mori Dream Space X . To achieve this, write the Cox ring $\mathbb{k}[x_1, \dots, x_d]$ of the toric variety \widetilde{X} as the semigroup algebra of the semigroup \mathbb{N}^d of effective torus-invariant Weil divisors in \widetilde{X} . Define the *label* of an arrow $a \in Q_1$, denoted $\text{div}(a) \in \mathbb{N}^d$, to be the divisor of zeroes in \widetilde{X} of the defining torus-invariant section $s \in H^0(\widetilde{X}, (E_{h(a)} \otimes E_{t(a)}^\vee)^{\vee\vee})$. More generally, the label of any path p in Q is the torus-invariant divisor $\text{div}(p) := \sum_{a \in \text{supp}(p)} \text{div}(a)$. It is often convenient to consider the corresponding labelling monomial

$$x^{\text{div}(p)} := \prod_{a \in \text{supp}(p)} x^{\text{div}(a)} \in \mathbb{k}[x_1, \dots, x_d]$$

in the Cox ring. Recall from (2.1) that our chosen presentation of the Cox ring of X determines an ideal $I_X \subset \mathbb{k}[x_1, \dots, x_d]$.

Definition 2.5. Consider the two-sided ideal

$$J_{\mathcal{L}} := \left(\sum_{p \in \Gamma} c_p p \in \mathbb{k}Q \mid \begin{array}{l} \exists \text{ finite set of paths } \Gamma \text{ in } Q \text{ that all have the same head and the} \\ \text{same tail, and } \exists \{c_p \in \mathbb{k} : p \in \Gamma\} \text{ such that } \sum_{p \in \Gamma} c_p x^{\text{div}(p)} \in I_X \end{array} \right)$$

in the path algebra $\mathbb{k}Q$. The pair $(Q, J_{\mathcal{L}})$ is the *bound quiver of sections* of the collection \mathcal{L} .

Proposition 2.6. *The quotient algebra $\mathbb{k}Q/J_{\mathcal{L}}$ is isomorphic to $\text{End}_{\mathcal{O}_X}(\bigoplus_{i \in Q_0} L_i)$, and each vertex $i \in Q_0$ satisfies $e_i(\mathbb{k}Q/J_{\mathcal{L}})e_0 \cong H^0(X, L_i)$.*

Proof. Adapt the proof of [6, Proposition 3.3] slightly, replacing $\text{Hom}(L_i, L_j)$ by $H^0(\widetilde{X}, (E_j \otimes E_i^\vee)^{\vee\vee})$ throughout, to see that the \mathbb{k} -algebra epimorphism

$$\widetilde{\eta}: \mathbb{k}Q \rightarrow \bigoplus_{i, j \in Q_0} H^0(\widetilde{X}, (E_j \otimes E_i^\vee)^{\vee\vee})$$

sending $\sum c_p p$ to $\sum c_p x^{\text{div}(p)}$ has kernel the following T -homogeneous ideal in $\mathbb{k}Q$:

$$J_{\widetilde{\mathcal{L}}} := (p^+ - p^- \in \mathbb{k}Q \mid h(p^+) = h(p^-), t(p^+) = t(p^-), \text{div}(p^+) = \text{div}(p^-))$$

For each $i, j \in Q_0$, isomorphism (2.4) implies that the restriction of the \mathbb{Z}^ρ -graded homomorphism τ from (2.2) defines a surjective \mathbb{k} -linear map $\tau_{ij}: H^0(\tilde{X}, (E_j \otimes E_i^\vee)^{\vee\vee}) \rightarrow H^0(X, L_j \otimes L_i^{-1})$ with kernel the \mathbb{k} -vector subspace $I_X \cap H^0(\tilde{X}, (E_j \otimes E_i^\vee)^{\vee\vee})$. The direct sum of all such maps defines the surjective, right-hand vertical map $\hat{\tau}$ in the commutative diagram of \mathbb{k} -algebras

$$(2.7) \quad \begin{array}{ccc} \mathbb{k}Q & \xrightarrow{\tilde{\eta}} & \bigoplus_{i,j \in Q_0} H^0(\tilde{X}, (E_j \otimes E_i^\vee)^{\vee\vee}) \\ \parallel & & \downarrow \hat{\tau} \\ \mathbb{k}Q & \xrightarrow{\eta} & \text{End}_{\mathcal{O}_X}(\bigoplus_{i \in Q_0} L_i) \end{array}$$

where the epimorphism η sends $\sum c_p p$ to the class $\sum c_p x^{\text{div}(p)} \bmod I_X$. The ideal $J_{\mathcal{L}}$ is the kernel of η , so the first statement holds. The second statement follows from the first since we have $L_0 = \mathcal{O}_X$ and we compose arrows and maps from right to left. \square

3. MORPHISM TO THE MULTIGRADED LINEAR SERIES

In this section we use the quiver of sections of a collection \mathcal{L} of line bundles on a Mori Dream Space X to define the corresponding multigraded linear series $|\mathcal{L}|$. Evaluating sections of line bundles defines a natural map from X to $|\mathcal{L}|$, and we establish necessary and sufficient conditions that make it a morphism and a closed immersion.

3.1. Toric quiver flag varieties. Let Q be a finite, connected quiver. A representation of Q consists of a \mathbb{k} -vector space W_i for $i \in Q_0$ and a \mathbb{k} -linear map $w_a: W_{t(a)} \rightarrow W_{h(a)}$ for $a \in Q_1$. It is convenient to write W as shorthand for $((W_i)_{i \in Q_0}, (w_a)_{a \in Q_1})$. The dimension vector of W is the vector $\underline{r} \in \mathbb{Z}^{Q_0}$ with components $r_i = \dim_{\mathbb{k}}(W_i)$ for $i \in Q_0$. A map of representations $\psi: W \rightarrow W'$ is a family $\psi_i: W_i \rightarrow W'_i$ of \mathbb{k} -linear maps for $i \in Q_0$ satisfying $w'_a \psi_{t(a)} = \psi_{h(a)} w_a$ for $a \in Q_1$. With composition defined componentwise, we obtain the abelian category of finite dimensional representations of Q . For $\theta \in \mathbb{Z}^{Q_0}$, define $\theta(W) := \sum_{0 \leq i \leq \rho} \theta_i \dim_{\mathbb{k}}(W_i)$. Following King [12], a representation W of Q is θ -semistable if $\theta(W) = 0$ and every subrepresentation $W' \subset W$ satisfies $\theta(W') \geq 0$. Moreover, W is θ -stable if the only subrepresentations W' with $\theta(W') = 0$ are 0 and W .

The incidence map $\text{inc}: \mathbb{Z}^{Q_1} \rightarrow \mathbb{Z}^{Q_0}$ defined by setting $\text{inc}(\chi_a) = \chi_{h(a)} - \chi_{t(a)}$ has image equal to the sublattice $\text{Wt}(Q) \subset \mathbb{Z}^{Q_0}$ of functions $\theta: Q_0 \rightarrow \mathbb{Z}$ satisfying $\sum_{i \in Q_0} \theta_i = 0$. The vectors $\{\chi_i - \chi_0 : i \neq 0\}$ form a \mathbb{Z} -basis for $\text{Wt}(Q)$. The $\text{Wt}(Q)$ -grading of $\mathbb{k}[y_a : a \in Q_1]$ determined by sending y_a to $\text{inc}(\chi_a)$ for $a \in Q_1$ induces a faithful action of the algebraic torus $G := \text{Hom}(\text{Wt}(Q), \mathbb{k}^\times)$ on $\mathbb{A}_{\mathbb{k}}^{Q_1} = \text{Spec } \mathbb{k}[y_a : a \in Q_1]$ in which $g = (g_i)_{i \in Q_0}$ acts on $w = (w_a)_{a \in Q_1}$ as $(g \cdot w)_a = g_{h(a)} w_a g_{t(a)}^{-1}$. For $\theta \in \text{Wt}(Q)$, let $\mathbb{k}[y_a : a \in Q_1]_\theta$ denote the θ -graded piece and

$$\mathbb{A}_{\mathbb{k}}^{Q_1} //_\theta G = \text{Proj} \left(\bigoplus_{j \geq 0} \mathbb{k}[y_a : a \in Q_1]_{j\theta} \right)$$

the categorical quotient of the open subset of θ -semistable points in $\mathbb{A}_{\mathbb{k}}^{Q_1}$.

Assume in addition that Q is acyclic with a unique source $0 \in Q_0$. The *toric quiver flag variety* Y_Q is the GIT quotient $\mathbb{A}_{\mathbb{k}}^{Q_1} //_\vartheta G$ linearised by the special weight $\vartheta := \sum_{i \in Q_0} (\chi_i - \chi_0) \in \text{Wt}(Q)$. Such varieties, studied initially by Craw–Smith [6] and in greater generality by Craw [5], can be characterised as follows:

Proposition 3.1. *Let Q be a finite, connected, acyclic quiver with a unique source $0 \in Q_0$ and special weight $\vartheta = \sum_{i \in Q_0} (\chi_i - \chi_0)$. The toric quiver flag variety Y_Q coincides with:*

- (i) *the GIT quotient $\mathbb{A}_{\mathbb{k}}^{Q_1} //_{\vartheta} G$ linearised by $\vartheta \in \text{Wt}(Q)$;*
- (ii) *the geometric quotient of $\mathbb{A}_{\mathbb{k}}^{Q_1} \setminus \mathbb{V}(B_Q)$ by the action of G , where the irrelevant ideal is*

$$B_Q := \left(\prod_{a \in \mathcal{T}} y_a : \mathcal{T} \text{ is a spanning tree of } Q \text{ rooted at } 0 \right) = \bigcap_{i \in Q_0 \setminus \{0\}} (y_a : h(a) = i);$$

- (iii) *the fine moduli space $\mathcal{M}_{\vartheta}(Q)$ of ϑ -stable representations of the quiver Q of dimension vector $\underline{r} = (1, \dots, 1) \in \mathbb{Z}^{Q_0}$.*

Moreover, Y_Q is a smooth projective toric variety obtained as a tower of projective space bundles.

Proof. See Craw–Smith [6, Proposition 3.8] and Craw [5, Theorem 3.3]. \square

The description of $Y_Q = \mathcal{M}_{\vartheta}(Q)$ as a fine moduli space of representations ensures that it carries a collection of tautological line bundles $\{\mathcal{W}_i : i \in Q_0\}$ with $\mathcal{W}_0 \cong \mathcal{O}_{Y_Q}$ and sheaf homomorphisms $\{\mathcal{W}_{t(a)} \rightarrow \mathcal{W}_{h(a)} : a \in Q_1\}$ whose restriction to the fibre over $\mathcal{M}_{\vartheta}(Q)$ encodes the corresponding representation $\{W_{t(a)} \rightarrow W_{h(a)} : a \in Q_1\}$. Moreover, the homomorphism of abelian groups $\text{Wt}(Q) \rightarrow \text{Pic}(Y_Q)$ satisfying

$$(3.1) \quad (\theta_0, \dots, \theta_r) \mapsto \mathcal{W}_1^{\theta_1} \otimes \dots \otimes \mathcal{W}_r^{\theta_r}$$

is an isomorphism. For more details, see [5, Sections 2-3].

3.2. Multigraded linear series. Let $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ be a collection of distinct, effective line bundles on a Mori Dream Space X . Lemma 2.4 guarantees that the corresponding quiver of sections Q is finite, connected, acyclic and has a unique source $0 \in Q_0$.

Definition 3.2. The *multigraded linear series* for \mathcal{L} is the toric quiver flag variety $|\mathcal{L}| := Y_Q$ of Q from Proposition 3.1. It carries tautological line bundles $\{\mathcal{W}_i : i \in Q_0\}$ with $\mathcal{W}_0 \cong \mathcal{O}_{Y_Q}$.

Remark 3.3. Just as Q is not precisely the quiver of sections of \mathcal{L} (see Remark 2.3), it is perhaps an abuse of terminology to call Y_Q the multigraded linear series of \mathcal{L} . Indeed, for the special case $\mathcal{L} = (\mathcal{O}_X, L_1)$ we have that $Y_Q \cong \mathbb{P}^*(H^0(E_1))$ is a projective space, but it need not coincide with the classical linear series $|L_1|$ because the epimorphism $\tau|_{H^0(\tilde{X}, E_1)} : H^0(\tilde{X}, E_1) \rightarrow H^0(X, L_1)$ from (2.2) need not be an isomorphism.

In order to study morphisms from X to the multigraded linear series $|\mathcal{L}|$, define

$$\tilde{\Phi} : \mathbb{k}[y_a : a \in Q_1] \rightarrow \mathbb{k}[x_1, \dots, x_d]$$

to be the \mathbb{k} -algebra homomorphism sending y_a to $x^{\text{div}(a)}$ for $a \in Q_1$. The actions of the groups $G = \text{Hom}(\text{Wt}(Q), \mathbb{k}^*)$ and $T = \text{Hom}(\mathbb{Z}^{\rho}, \mathbb{k}^*)$ on $\mathbb{k}[y_a : a \in Q_1]$ and $\mathbb{k}[x_1, \dots, x_d]$ respectively arise from the horizontal semigroup homomorphisms in the diagram

$$(3.2) \quad \begin{array}{ccc} \mathbb{N}^{Q_1} & \xrightarrow{\text{inc}} & \text{Wt}(Q) \\ \text{div} \downarrow & & \downarrow \text{pic} \\ \mathbb{N}^d & \xrightarrow[\mathfrak{g}]{\text{deg}} & \mathbb{Z}^{\rho} \end{array}$$

where the vertical maps satisfy $\operatorname{div}(\chi_a) = \operatorname{div}(a)$ for $a \in Q_1$ and $\operatorname{pic}(\chi_i) = E_i$ for $i \in Q_0$. The map $\tilde{\Phi}$ is equivariant with respect to these actions precisely because (3.2) commutes. Under the identification of $\operatorname{Wt}(Q)$ with the Picard group of $|\mathcal{L}|$, the subspace of the Cox ring $\mathbb{k}[y_a : a \in Q_1]$ of $|\mathcal{L}|$ spanned by monomials of weight $\theta \in \operatorname{Wt}(Q)$ coincides with $H^0(\mathcal{W}_1^{\theta_1} \otimes \cdots \otimes \mathcal{W}_r^{\theta_r})$.

Since the T -action on $\operatorname{Cox}(X)$ is compatible with that on $\mathbb{k}[x_1, \dots, x_d]$ by (2.2), the map

$$\Phi := \tau \circ \tilde{\Phi}: \mathbb{k}[y_a : a \in Q_1] \longrightarrow \operatorname{Cox}(X)$$

is equivariant. The induced equivariant morphism $\Phi^*: \mathbb{V}(I_X) \rightarrow \mathbb{A}_{\mathbb{k}}^{Q_1}$ descends to a rational map $\varphi_{|\mathcal{L}|}: X \dashrightarrow |\mathcal{L}|$.

Proposition 3.4. *Let $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ be a collection of distinct, effective line bundles on X . The rational map $\varphi_{|\mathcal{L}|}: X \dashrightarrow |\mathcal{L}|$ is a morphism if and only if L_i is basepoint-free for $1 \leq i \leq r$.*

Proof. For $x \in X$ choose any lift $\tilde{x} \in \mathbb{V}(I_X) \setminus \mathbb{V}(B_X)$. The G -orbit of the quiver representation $\Phi^*(\tilde{x}) \in \mathbb{A}_{\mathbb{k}}^{Q_1}$, which is independent of the choice of lift, is obtained by evaluating the labels on arrows at \tilde{x} , that is, by evaluating sections of the bundles $L_{h(a)} \otimes L_{t(a)}^{-1}$ at x . The rational map $\varphi_{|\mathcal{L}|}: X \dashrightarrow |\mathcal{L}|$ is a morphism if and only if every such $\Phi^*(\tilde{x}) \in \mathbb{A}_{\mathbb{k}}^{Q_1}$ is ϑ -stable. Let $W' = ((W'_i)_{i \in Q_0}, (w'_a)_{a \in Q_1})$ be a proper subrepresentation of $\Phi^*(\tilde{x})$. Since $\vartheta_0 = -r$ and $\vartheta_i > 0$ for $i > 0$, the submodule W' of $\Phi^*(\tilde{x})$ is ϑ -destabilising if and only if $\dim_{\mathbb{k}}(W'_0) = 1$ and there exists $i > 0$ such that for every path $p = a_\ell \cdots a_1$ from 0 to i , the composition $w'_{a_\ell} \cdots w'_{a_1}$ is the zero map. In particular, $\Phi^*(\tilde{x}) \in \mathbb{A}_{\mathbb{k}}^{Q_1}$ is ϑ -unstable if and only if there exists $i > 0$ such that the evaluation of every section of L_i at x equals zero. Equivalently, $\Phi^*(\tilde{x}) \in \mathbb{A}_{\mathbb{k}}^{Q_1}$ is ϑ -stable if and only if L_i is basepoint-free for $1 \leq i \leq r$. \square

The Cox ring of X is a unique factorisation domain, so $\ker(\Phi)$ is prime and hence so is the ideal

$$(3.3) \quad I_Q := \left(f \in \mathbb{k}[y_a : a \in Q_1] : f \in \ker(\Phi) \text{ is } \operatorname{Wt}(Q)\text{-homogeneous} \right)$$

generated by its $\operatorname{Wt}(Q)$ -homogeneous elements. This ideal can be computed explicitly as the kernel of the \mathbb{k} -algebra homomorphism

$$(3.4) \quad \Psi: \mathbb{k}[y_a : a \in Q_1] \rightarrow \operatorname{Cox}(X) \otimes_{\mathbb{k}} \mathbb{k}[\operatorname{Wt}(Q)]$$

satisfying $\Psi(y_a) = t_{h(a)} x^{\operatorname{div}(a)} t_{t(a)}^{-1}$ for $a \in Q_1$; see Winn [15, Chapter 5] for details. This ideal cuts out the image of the morphism constructed in Proposition 3.4 as follows.

Proposition 3.5. *Let $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ be a collection of distinct, basepoint-free line bundles on X with quiver of sections Q . Then*

- (i) *the image of the morphism $\varphi_{|\mathcal{L}|}: X \rightarrow |\mathcal{L}|$ is $\mathbb{V}(I_Q) //_{\vartheta} G$; and*
- (ii) *the tautological line bundles on $|\mathcal{L}|$ satisfy $\varphi_{|\mathcal{L}|}^*(\mathcal{W}_i) = L_i$ for $i \in Q_0$.*

Proof. Since X is complete, the image of $\varphi_{|\mathcal{L}|}$ is a closed subscheme of $|\mathcal{L}|$. The geometric quotient construction of $|\mathcal{L}|$ from Proposition 3.1(i) implies that the image is therefore the geometric quotient of a G -invariant closed subscheme of $\mathbb{A}_{\mathbb{k}}^{Q_1} \setminus \mathbb{V}(B_Q)$. The affine variety $\mathbb{V}(\ker(\Phi))$ is the image of the equivariant morphism $\operatorname{Spec}(\operatorname{Cox}(X)) \rightarrow \mathbb{A}_{\mathbb{k}}^{Q_1}$ induced by Φ , and the variety $\mathbb{V}(I_Q)$ cut out by the $\operatorname{Wt}(Q)$ -homogeneous part of $\ker(\Phi)$ is the minimal G -invariant algebraic

set in \mathbb{A}^{Q_1} containing all G -orbits intersecting $\mathbb{V}(\ker(\Phi))$. The image of $\varphi_{|\mathcal{L}|}$ is therefore the geometric quotient of $\mathbb{V}(I_Q) \setminus \mathbb{V}(B_Q)$ by the action of G . This coincides with the GIT quotient $\mathbb{V}(I_Q) //_{\vartheta} G$ by Proposition 3.1, so (i) holds. For part (ii), the tautological bundle \mathcal{W}_i on $|\mathcal{L}|$ corresponds to the weight $\chi_i - \chi_0 \in \text{Wt}(Q)$ under the isomorphism (3.1). Since the equivariant morphism $\text{Spec}(\text{Cox}(X)) \rightarrow \mathbb{A}_{\mathbb{k}}^{Q_1}$ factors through $\mathbb{A}_{\mathbb{k}}^d$, examining (2.2) and diagram (3.2) shows that $\varphi_{|\mathcal{L}|}^*(\mathcal{W}_i) = (\psi \circ \text{pic})(\chi_i - \chi_0) = \psi(E_i) = L_i$ for $i \in Q_0$. \square

Proof of Theorem 1.1. Proposition 3.4 establishes that $\varphi_{|\mathcal{L}|}: X \dashrightarrow |\mathcal{L}|$ is a morphism if and only if L_i is basepoint-free for $1 \leq i \leq r$. Proposition 3.5 then presents the image explicitly as a geometric quotient, and establishes that the tautological line bundles on $|\mathcal{L}|$ satisfy $\varphi_{|\mathcal{L}|}^*(\mathcal{W}_i) = L_i$ for $i \in Q_0$ as required. \square

Remark 3.6. The list of reflexive sheaves $\widetilde{\mathcal{L}}$ on \widetilde{X} determines the ideal

$$(3.5) \quad I_{\widetilde{Q}} = \left(f \in \mathbb{k}[y_a : a \in Q_1] : f \in \ker(\widetilde{\Phi}) \text{ is } \text{Wt}(Q)\text{-homogeneous} \right)$$

obtained as the toric ideal of the semigroup homomorphism $\text{inc} \oplus \text{div}: \mathbb{N}^{Q_1} \rightarrow \text{Wt}(Q) \oplus \mathbb{N}^d$. If each reflexive sheaf in $\widetilde{\mathcal{L}}$ is a basepoint-free line bundle on \widetilde{X} , then [6, Theorem 1] gives a morphism $\varphi_{|\widetilde{\mathcal{L}}|}: \widetilde{X} \rightarrow \mathbb{V}(I_{\widetilde{Q}}) //_{\vartheta} G$ whose restriction to X is the morphism $\varphi_{|\mathcal{L}|}: X \rightarrow \mathbb{V}(I_Q) //_{\vartheta} G$ from Proposition 3.5. However, this is typically not the case as in Example 3.10.

3.3. Criteria for closed immersion. A collection \mathcal{L} is said to be *very ample* if the morphism $\varphi_{|\mathcal{L}|}$ from Proposition 3.4 is a closed immersion. We now introduce a necessary and sufficient condition for \mathcal{L} to be very ample. We (enhance and) adapt the proofs of [5, Proposition 5.7] and [6, Corollary 4.10] to our situation because Q is not precisely the quiver of sections for \mathcal{L} (see Remarks 2.3 and 3.3).

Theorem 3.7. *Let $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ be a collection of distinct, basepoint-free line bundles on X . The following are equivalent:*

- (i) *the morphism $\varphi_{|\mathcal{L}|}: X \rightarrow |\mathcal{L}|$ is a closed immersion;*
- (ii) *the image of the multiplication map*

$$(3.6) \quad H^0(L_1) \otimes \dots \otimes H^0(L_r) \longrightarrow H^0(L_1 \otimes \dots \otimes L_r).$$

is a very ample linear series;

- (iii) *the map $\prod_{1 \leq i \leq r} \varphi_{|L_i|}: X \rightarrow |L_1| \times \dots \times |L_r|$ is a closed immersion.*

Proof. The toric variety $|\mathcal{L}|$ is smooth, so the ample bundle $\vartheta = \mathcal{W}_1 \otimes \dots \otimes \mathcal{W}_r$ determines the closed immersion $\varphi_{|\vartheta|}: |\mathcal{L}| \rightarrow \mathbb{P}^*(H^0(\vartheta))$. The composition $\varphi_{|\vartheta|} \circ \varphi_{|\mathcal{L}|}: X \rightarrow \mathbb{P}^*(H^0(\vartheta))$ is determined by the line bundle $(\varphi_{|\vartheta|} \circ \varphi_{|\mathcal{L}|})^*(\vartheta) = (\psi \circ \text{pic})(\theta) = L_1 \otimes \dots \otimes L_r$ and the subspace of sections $\Phi(H^0(\vartheta)) \subseteq H^0(L_1 \otimes \dots \otimes L_r)$. We claim that $\Phi(H^0(\vartheta))$ coincides with the image V of the multiplication map (3.6), in which case $\varphi_{|\vartheta|} \circ \varphi_{|\mathcal{L}|}$ coincides with the (a priori rational) map $\varphi_V: X \rightarrow \mathbb{P}^*(V)$ to the classical linear series. Indeed, for $\theta = (\theta_0, \dots, \theta_r) \in \text{Wt}(Q)$, the restriction of Φ to the subspace spanned by monomials of weight θ defines a \mathbb{k} -linear map

$$\Phi_{\theta}: H^0(\mathcal{W}_1^{\theta_1} \otimes \dots \otimes \mathcal{W}_r^{\theta_r}) \rightarrow H^0(L_1^{\theta_1} \otimes \dots \otimes L_r^{\theta_r})$$

because $(\psi \circ \text{pic})(\theta) = L_1^{\theta_1} \otimes \cdots \otimes L_r^{\theta_r}$. In particular, the map Φ_{ϑ} for $\vartheta = \sum_{1 \leq i \leq r} (\chi_i - \chi_0)$ and the product $\otimes_{1 \leq i \leq r} \Phi_{(\chi_i - \chi_0)}$ fit in to a commutative diagram of \mathbb{k} -vector spaces

$$(3.7) \quad \begin{array}{ccc} H^0(\mathcal{W}_1) \otimes \cdots \otimes H^0(\mathcal{W}_r) & \longrightarrow & H^0(\mathcal{W}_1 \otimes \cdots \otimes \mathcal{W}_r) \\ \otimes_{1 \leq i \leq r} \Phi_{(\chi_i - \chi_0)} \downarrow & & \downarrow \Phi_{\vartheta} \\ H^0(L_1) \otimes \cdots \otimes H^0(L_r) & \longrightarrow & H^0(L_1 \otimes \cdots \otimes L_r) \end{array}$$

in which the horizontal maps are given by multiplication. For $1 \leq i \leq r$, the map $\Phi_{(\chi_i - \chi_0)}$ can be obtained by composing three surjective maps, namely, the isomorphism $H^0(\mathcal{W}_i) \rightarrow e_i(\mathbb{k}Q)e_0$ from [5, Corollary 3.5], the restriction to $e_i(\mathbb{k}Q)e_0$ of the epimorphism $\tilde{\eta}: \mathbb{k}Q \rightarrow \text{End}(\bigoplus_{i \in Q_0} E_i)$ from the proof of Proposition 2.6, and the restricted map $\tau|_{H^0(E_i)}: H^0(E_i) \rightarrow H^0(L_i)$ from (2.2). It follows that $\bigotimes_{1 \leq i \leq r} \Phi_{(\chi_i - \chi_0)}$ is surjective. Every monomial of weight ϑ in $\mathbb{k}[y_a : a \in Q_1]$ can be decomposed as a product of monomials of weight $\chi_i - \chi_0 \in \text{Wt}(Q)$ for $i \in Q_0$, so the top horizontal map in diagram (3.7) is surjective. Commutativity of the diagram then implies that the image of Φ_{ϑ} coincides with the image V of (3.6). This proves the claim.

Since V is the image of the multiplication map (3.6), the morphism $\varphi_V: X \rightarrow \mathbb{P}^*(V)$ is the composition of the product $\prod_{1 \leq i \leq r} \varphi_{|L_i|}: X \rightarrow |L_1| \times \cdots \times |L_r|$ of morphisms to the classical linear series and the appropriate Segre embedding to $\mathbb{P}^*(V)$. The claim implies that the diagram

$$\begin{array}{ccccc} |L_1| \times \cdots \times |L_r| & \xrightarrow{\text{Segre}} & \mathbb{P}^*(V) & \xrightarrow{\iota} & \mathbb{P}^*(H^0(\vartheta)) \\ \uparrow \prod_{1 \leq i \leq r} \varphi_{|L_i|} & & & & \uparrow \varphi_{|\vartheta|} \\ X & \xrightarrow{\varphi_{|\mathcal{L}|}} & |\mathcal{L}| & & \end{array}$$

commutes, where ι is the closed immersion of projective spaces induced by Φ_{ϑ} . Three maps in the diagram are closed immersions, so $\varphi_{|\mathcal{L}|}$ is a closed immersion if and only if $\prod_{1 \leq i \leq r} \varphi_{|L_i|}$ is a closed immersion if and only if the linear series V is very ample as required. \square

Remark 3.8. Neither of the maps from statements (i) and (iii) of Theorem 3.7 factors through the other. Typically $|\mathcal{L}|$ has much lower dimension than $|L_1| \times \cdots \times |L_r|$, so the multigraded linear series is a more efficient multigraded ambient space than the product.

Corollary 3.9. *Let L_1, \dots, L_{r-1} be distinct, basepoint-free line bundles on X . If the subsemigroup of $\text{Pic}(X)$ generated by L_1, \dots, L_{r-1} contains an ample bundle, then there exists a line bundle L_r such that the quiver of sections for $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ is very ample.*

Proof. Theorem 3.7 implies that $\varphi_{|\mathcal{L}|}$ is a closed immersion if $L_1 \otimes \cdots \otimes L_r$ is very ample and the map (3.6) is surjective. The proof of [6, Proposition 4.14] now applies verbatim. \square

Example 3.10. Continuing Example 2.1, let X_4 be the del Pezzo surface for which the ample linearisation $\chi = 11H - 5R_1 - 3R_2 - 2R_3 - R_4$ defines $\widetilde{X}_4 := \mathbb{A}^{10} //_{\chi} T$. We compute using the intersection pairing that each line bundle in the list

$$(3.8) \quad \mathcal{L} = (\mathcal{O}_{X_4}, H, 2H - R_1, 2H - R_2, 2H - R_3, 2H - R_4, 2H)$$

is basepoint-free but not ample. Write $\widetilde{\mathcal{L}} = (E_0, E_1, \dots, E_6)$. Since each E_i is basepoint-free on some ambient toric variety, the code from [13, Example 2.11] computes the irrelevant ideal for the GIT quotient $\mathbb{A}_{\mathbb{k}}^d //_{E_i} T$ determined by the corresponding linearisation $E_i \in \mathbb{Z}^5$. By comparing

each with the irrelevant ideal of $\chi \in \mathbb{Z}^5$ we see that E_3, E_4, E_5 are not basepoint-free line bundles on \widetilde{X}_4 . In particular, we cannot deduce that $\varphi_{|\mathcal{L}|}$ is a morphism simply by restriction from the toric case (compare Remark 3.6), though it is a morphism by Proposition 3.4.

To investigate $\varphi_{|\mathcal{L}|}$ directly in this case, the quiver of sections Q is shown in Figure 1, where each arrow is labelled by the torus-invariant section of the relevant reflexive sheaf on \widetilde{X}_4 . Arrows




FIGURE 1. A quiver of sections for a collection on X_4

with tail at 0 are listed a_1, \dots, a_6 from the top of Figure 1 to the bottom; list those with tail at 1 as a_7, \dots, a_{18} from the top of the figure to the bottom; and list those with head at 6 as a_{19}, \dots, a_{22} from the top to the bottom. Likewise, list the coordinates of $\mathbb{A}_{\mathbb{k}}^{Q_1}$ as y_1, \dots, y_{22} , and compute the kernel of (3.4) to obtain the ideal

$$I_Q = \left(\begin{array}{c} y_{16} - y_{17} + y_{18}, y_{13} - y_{14} + y_{15}, y_{10} - y_{11} + y_{12}, y_7 - y_8 + y_9, y_3 - y_5 + y_6 \\ y_2 - y_4 + y_6, y_1 - y_4 + y_5, y_{15}y_{21} - y_{18}y_{22}, y_{12}y_{20} - y_{17}y_{22}, y_{11}y_{20} - y_{14}y_{21} \\ y_9y_{19} - y_{17}y_{22} + y_{18}y_{22}, y_8y_{19} - y_{14}y_{21} + y_{18}y_{22}, y_6y_{17} - y_5y_{18}, y_6y_{14} - y_4y_{15} \\ y_5y_{11} - y_4y_{12}, y_5y_8 - y_6y_8 - y_4y_9 + y_6y_9, y_8y_{15}y_{17} - y_9y_{14}y_{18} - y_8y_{15}y_{18} + y_9y_{15}y_{18} \\ y_{11}y_{15}y_{17} - y_{12}y_{14}y_{18}, y_9y_{11}y_{17} - y_8y_{12}y_{17} + y_8y_{12}y_{18} - y_9y_{12}y_{18} \\ y_9y_{11}y_{14} - y_8y_{12}y_{14} + y_8y_{11}y_{15} - y_9y_{11}y_{15} \end{array} \right)$$

that cuts out the image of $\varphi_{|\mathcal{L}|}: X_4 \rightarrow |\mathcal{L}|$.

We claim that $\varphi_{|\mathcal{L}|}$ is a closed immersion, and hence $X_4 \cong \mathbb{V}(I_Q)_{/\vartheta} G$. Indeed, for $1 \leq i \leq 4$ we have $L_{i+1} = 2H - R_i$, and the intersection pairing shows that $\varphi_{|L_{i+1}|}: X_4 \rightarrow \mathbb{P}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(1))$ contracts the (-1) -curves $\{R_j : j \neq i\}$ but not R_i . A simple case-by-case analysis shows that the morphism $\prod_{2 \leq i \leq 5} \varphi_{|L_i|}$ separates all points and tangent vectors of X_4 and hence so does $\prod_{1 \leq i \leq 6} \varphi_{|L_i|}$. We deduce from Theorem 3.7 that $\varphi_{|\mathcal{L}|}: X_4 \rightarrow |\mathcal{L}|$ is a closed immersion.

4. FINE MODULI OF BOUND QUIVER REPRESENTATIONS

This section establishes when the morphism $\varphi_{|\mathcal{L}|}: X \rightarrow |\mathcal{L}|$ induces an isomorphism between the Mori Dream Space X and a fine moduli space $\mathcal{M}_{\vartheta}(\text{mod-}A)$ of ϑ -stable modules over the algebra $A_{\mathcal{L}} = \text{End}(\bigoplus_{i \in Q_0} L_i)$. Our main algebraic result is an efficient construction for collections of line bundles that induce the isomorphism.

4.1. Bound quiver representations. Let Q be a quiver. For any nontrivial path $p = a_k \cdots a_1$ in Q , define the monomial $y_p := y_{a_k} \cdots y_{a_1} \in \mathbb{k}[y_a : a \in Q_1]$, and for any representation W of Q , define $w_p: W_{t(p)} \rightarrow W_{h(p)}$ to be the \mathbb{k} -linear map $w_p = w_{a_k} \cdots w_{a_1}$ obtained by composition.

Let $J \subset \mathbb{k}Q$ be a two-sided ideal of relations with generators of the form $\sum_{p \in \Gamma} c_p p$, where each Γ is a finite set of paths that share the same head and the same tail. A representation W of Q is a representation of the bound quiver (Q, J) if and only if $\sum_{p \in \Gamma} c_p w_p = 0$ for each Γ arising in the definition of J . A point in representation space $(w_a) \in \mathbb{A}_{\mathbb{k}}^{Q_1}$ defines a representation of (Q, J) if and only if it lies in the subscheme $\mathbb{V}(I_J)$ cut out by the ideal

$$I_J := \left(\sum_{p \in \Gamma} c_p y_p \in \mathbb{k}[y_a : a \in Q_1] \mid \sum_{p \in \Gamma} c_p p \in J \right)$$

of relations in $\mathbb{k}[y_a : a \in Q_1]$. The ideal I_J is $\text{Wt}(Q)$ -homogeneous, so $\mathbb{V}(I_J)$ is G -invariant and the GIT quotient

$$(4.1) \quad \mathcal{M}_{\vartheta}(Q, J) := \mathbb{V}(I_J) //_{\vartheta} G = \text{Proj} \left(\bigoplus_{j \geq 0} (\mathbb{k}[y_a : a \in Q_1] / I_J)_{j\vartheta} \right)$$

is the fine moduli space of ϑ -stable representations of (Q, J) with dimension vector $(1, \dots, 1)$. The tautological bundles on $\mathcal{M}_{\vartheta}(Q, J)$ are obtained from those on $\mathcal{M}_{\vartheta}(Q)$ by restriction.

Remark 4.1. The abelian category of finite-dimensional representations of (Q, J) is equivalent to the category of finitely-generated $A := \mathbb{k}Q/J$ -modules, so $\mathcal{M}_{\vartheta}(Q, J)$ is equivalently the fine moduli space of ϑ -stable A -modules that are isomorphic as $(\bigoplus_{i \in Q_0} \mathbb{k}e_i)$ -modules to $\bigoplus_{i \in Q_0} \mathbb{k}e_i$.

4.2. Ideals of relations. A list \mathcal{L} of line bundles on X defines a pair of two-sided ideals in $\mathbb{k}Q$ and hence a pair of ideals of relations in $\mathbb{k}[y_a : a \in Q_1]$. First, the ideal $J_{\mathcal{L}}$ from Definition 2.5 determines the ideal of relations $I_{\mathcal{L}} := I_{(J_{\mathcal{L}})}$, that is, the ideal

$$(4.2) \quad I_{\mathcal{L}} := \left(\sum_{p \in \Gamma} c_p y_p \in \mathbb{k}[y_a : a \in Q_1] \mid \begin{array}{l} \exists \text{ finite set of paths } \Gamma \text{ with same head and same} \\ \text{tail, and } \exists c_p \in \mathbb{k} \text{ such that } \tilde{\Phi}(\sum_{p \in \Gamma} c_p y_p) \in I_X \end{array} \right).$$

Each generator of $I_{\mathcal{L}}$ is $\text{Wt}(Q)$ -homogeneous and lies in $\ker(\Phi)$, so $I_{\mathcal{L}}$ is contained in the prime ideal I_Q from (3.3). Winn [15, Chapter 5] presents code that calculates $I_{\mathcal{L}}$ explicitly.

In addition, the kernel $J_{\tilde{\mathcal{L}}}$ of the epimorphism $\tilde{\eta}: \mathbb{k}Q \rightarrow \text{End}_{\mathcal{O}_{\tilde{X}}}(\bigoplus_{i \in Q_0} E_i)$ from the proof of Proposition 2.6 determines the noncommutative toric ideal of relations $I_{\tilde{\mathcal{L}}} := I_{(J_{\tilde{\mathcal{L}}})}$, where

$$(4.3) \quad I_{\tilde{\mathcal{L}}} := \left(\sum_{p \in \tilde{\Gamma}} c_p y_p \in \mathbb{k}[y_a : a \in Q_1] \mid \begin{array}{l} \exists \text{ finite set of paths } \tilde{\Gamma} \text{ with same head and same} \\ \text{tail, and } \exists c_p \in \mathbb{k} \text{ such that } \tilde{\Phi}(\sum_{p \in \tilde{\Gamma}} c_p y_p) = 0 \end{array} \right).$$

We have that $I_{\tilde{\mathcal{L}}}$ is contained both in the ideal of equations $I_{\tilde{Q}}$ from (3.5), and in $I_{\mathcal{L}}$ from (4.2). Compute the affine varieties in $\mathbb{A}_{\mathbb{k}}^{Q_1}$ cut out by the ideals $I_{\tilde{\mathcal{L}}}, I_{\tilde{Q}}, I_{\mathcal{L}}, I_Q \subset \mathbb{k}[y_a : a \in Q_1]$, remove from each the ϑ -unstable locus $\mathbb{V}(B_Q)$, and compute the geometric quotient by the action of G to obtain the left-hand square in the commutative diagram of GIT quotients

$$(4.4) \quad \begin{array}{ccccc} \mathbb{V}(I_{\tilde{Q}}) //_{\vartheta} G & \longrightarrow & \mathcal{M}_{\vartheta}(Q, J_{\tilde{\mathcal{L}}}) & \longrightarrow & \mathbb{A}_{\mathbb{k}}^{Q_1} //_{\vartheta} G \\ \uparrow & & \uparrow & & \parallel \\ \mathbb{V}(I_Q) //_{\vartheta} G & \longrightarrow & \mathcal{M}_{\vartheta}(Q, J_{\mathcal{L}}) & \longrightarrow & |\mathcal{L}| \end{array}$$

in which each morphism is a closed immersion.

Recall that for ideals $B, I \subseteq \mathbb{k}[y_a : a \in Q_1]$, the saturation of I by B is the ideal

$$(I : B^{\infty}) = (g \in \mathbb{k}[y_a : a \in Q_1] \mid h^N g \in I \text{ for some } h \in B_Q \text{ and } N > 0).$$

Theorem 4.2. *If \mathcal{L} is a collection of distinct, basepoint-free line bundles on X , then the induced morphism*

$$(4.5) \quad \varphi_{|\mathcal{L}|}: X \longrightarrow \mathcal{M}_\vartheta(Q, J_{\mathcal{L}})$$

is surjective if I_Q coincides with the saturation $(I_{\mathcal{L}} : B_Q^\infty)$. In particular, if \mathcal{L} is very ample and $I_Q = (I_{\mathcal{L}} : B_Q^\infty)$ then (4.5) is an isomorphism.

Proof. It suffices by Theorem 3.7 to show that the closed immersion $\mathbb{V}(I_Q) //_\vartheta G \rightarrow \mathbb{V}(I_{\mathcal{L}}) //_\vartheta G$ is an isomorphism. Proposition 3.1 shows that the ideal B_Q cuts out the ϑ -unstable locus in $\mathbb{A}_{\mathbb{k}}^{Q_1}$, so we need only show that $\mathbb{V}(I_Q) \setminus \mathbb{V}(B_Q)$ is isomorphic to $\mathbb{V}(I_{\mathcal{L}}) \setminus \mathbb{V}(B_Q)$. Since I_Q is prime, this holds if $I_Q = (I_{\mathcal{L}} : B_Q^\infty)$. The second statement is immediate. \square

Remark 4.3. In light of Proposition 2.6 and Remark 4.1, when the map (4.5) is an isomorphism then we describe the Mori Dream Space X as the fine moduli space $\mathcal{M}_\vartheta(\text{mod-}A)$ of ϑ -stable modules over $A := \text{End}(\bigoplus_{i \in Q_0} L_i)$ that are isomorphic as $(\bigoplus_{i \in Q_0} \mathbb{k}e_i)$ -modules to $\bigoplus_{i \in Q_0} \mathbb{k}e_i$.

4.3. Digression on multigraded regularity. The notion of multigraded regularity for a sheaf with respect to a collection of basepoint-free line bundles on a projective variety, introduced by MacLagan–Smith [14] in the toric case and Hering–Schenck–Smith [9] in general, provides a multigraded variant of Castelnuovo–Mumford regularity. The special case of \mathcal{O}_X -multigraded regularity provides the key technical tool in the proof of Theorem 1.2 (see Section 4.4 to follow), and for convenience we recall the definition and an important property.

Consider a list $\mathcal{E} = (E_1, \dots, E_\ell)$ of basepoint-free line bundles on a projective variety X . For an integer vector $\beta = (\beta_1, \dots, \beta_\ell) \in \mathbb{Z}^\ell$ we set $E^\beta = E_1^{\beta_1} \otimes \dots \otimes E_\ell^{\beta_\ell}$.

Definition 4.4. A line bundle L on X is \mathcal{O}_X -regular with respect to \mathcal{E} if $H^i(X, L \otimes E^{-\beta}) = 0$ for all $i > 0$ and for all $\beta \in \mathbb{N}^\ell$ with $\beta_1 + \dots + \beta_\ell = i$.

This notion was used implicitly (see [6, Proposition 4.14]) in the proof of Corollary 3.9, but it becomes essential in the following section. The crucial property for us is the following sufficient condition for the multiplication map on global sections to be surjective (see [9, Theorem 2.1(2)] [14, Theorem 6.9]):

Proposition 4.5 ([9, 14]). *Let F be a coherent sheaf on X . If F is \mathcal{O}_X -regular then the map*

$$H^0(X, F \otimes E^\alpha) \otimes_{\mathbb{k}} H^0(X, E^\beta) \longrightarrow H^0(X, F \otimes E^{\alpha+\beta})$$

is surjective for all $\alpha, \beta \in \mathbb{N}^\ell$.

4.4. Main algebraic result. We now work towards our main algebraic result which exhibits many collections of line bundles on X for which the morphism from (4.5) is an isomorphism, thereby providing a noncommutative algebraic construction of X as in Remark 4.3.

We first introduce the collections of interest. Choose generators $g_1, \dots, g_m \in \mathbb{k}[x_1, \dots, x_d]$ of the ideal I_X , set $\delta_0 := \max_{1 \leq j \leq m} \{\text{total degree of } g_j\}$ and define

$$(4.6) \quad \delta := \begin{cases} \delta_0/2 & \text{if } \delta_0 \text{ is even;} \\ (\delta_0 + 1)/2 & \text{otherwise.} \end{cases}$$

Consider line bundles L_1, \dots, L_{r-2} on X for which the corresponding rank one reflexive sheaves $E_1 = \psi^{-1}(L_1), \dots, E_{r-2} = \psi^{-1}(L_{r-2})$ on \tilde{X} are basepoint-free line bundles such that the

subsemigroup of $\text{Pic}(\tilde{X})$ generated by E_1, \dots, E_{r-2} contains an ample line bundle. Choose sufficiently positive integers $\beta_1, \dots, \beta_{r-2}$ to ensure that $E := E_1^{\beta_1} \otimes \dots \otimes E_{r-2}^{\beta_{r-2}}$ is both normally generated and $\mathcal{O}_{\tilde{X}}$ -regular with respect to E_1, \dots, E_{r-2} (see Definition 4.4) and, moreover, that $E^{2\delta}$ is very ample. Define $E_{r-1} := E^\delta$ and $E_r := E^{2\delta}$ (assuming that each is distinct from E_1, \dots, E_{r-2}). Augment the list L_1, \dots, L_{r-2} on X with $L_0 = \mathcal{O}_X$, $L_{r-1} := \psi(E_{r-1})$ and $L_r := \psi(E_r)$ to obtain a collection

$$(4.7) \quad \mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$$

of basepoint-free line bundles on X . Let Q denote the quiver of sections of \mathcal{L} . The corresponding collection of line bundles $\tilde{\mathcal{L}} := (\mathcal{O}_{\tilde{X}}, E_1, \dots, E_r)$ on \tilde{X} takes the form of the collections constructed in the proof of [6, Theorem 5.5], and hence by the proof of that result we have

$$(4.8) \quad I_{\tilde{Q}} = (I_{\tilde{\mathcal{L}}} : B_Q^\infty)$$

and the induced morphism $\varphi_{|\tilde{\mathcal{L}}|} : \tilde{X} \rightarrow \mathbb{A}^{Q_1} //_{\vartheta} G$ is a closed immersion with image $\mathbb{V}(I_{\tilde{Q}}) //_{\vartheta} G$ isomorphic to $\mathcal{M}_{\vartheta}(Q, J_{\tilde{\mathcal{L}}})$.

Remark 4.6. (1) It follows that each collection (4.7) determines a commutative diagram

$$(4.9) \quad \begin{array}{ccccccc} \tilde{X} & \xrightarrow{\varphi_{|\tilde{\mathcal{L}}|}} & \mathbb{V}(I_{\tilde{Q}}) //_{\vartheta} G & \xrightarrow{\cong} & \mathcal{M}_{\vartheta}(Q, J_{\tilde{\mathcal{L}}}) & \longrightarrow & \mathbb{A}_{\mathbb{k}}^{Q_1} //_{\vartheta} G \\ \uparrow & & \uparrow & & \uparrow & & \parallel \\ X & \xrightarrow{\varphi_{|\mathcal{L}|}} & \mathbb{V}(I_Q) //_{\vartheta} G & \longrightarrow & \mathcal{M}_{\vartheta}(Q, J_{\mathcal{L}}) & \longrightarrow & |\mathcal{L}| \end{array}$$

in which every morphism is a closed immersion and the indicated map is an isomorphism.

- (2) Since E is $\mathcal{O}_{\tilde{X}}$ -regular with respect to E_1, \dots, E_{r-2} , since each $\beta_i > 0$ and since $E_{r-1} = E_1^{\delta\beta_1} \otimes \dots \otimes E_{r-2}^{\delta\beta_{r-2}}$, Proposition 4.5 implies that the multiplication map

$$H^0(E_{r-1} \otimes E_i^{-1}) \otimes_{\mathbb{k}} H^0(E_{r-1}) \longrightarrow H^0(E_r \otimes E_i^{-1})$$

is surjective for $1 \leq i \leq r-1$. Since each E_i is invertible, it follows from (2.6) that every path in Q from 0 to r passes through $r-1$.

- (3) For clarity in what follows, we work with elements of $\mathbb{k}[y_a : a \in Q_1]$ modulo the relation \sim in which polynomials are equivalent when their difference lies in $I_{\tilde{Q}}$. Since $I_{\tilde{Q}}$ is the toric ideal of the semigroup homomorphism $\text{inc} \oplus \text{div} : \mathbb{N}^{Q_1} \rightarrow \text{Wt}(Q) \oplus \mathbb{N}^d$, monomials satisfy $y^{\mathbf{m}} \sim y^{\mathbf{m}'}$ if and only if $\text{inc}(\mathbf{m} - \mathbf{m}') = 0$ and $\text{div}(\mathbf{m} - \mathbf{m}') = 0$, that is, $y^{\mathbf{m}} \sim y^{\mathbf{m}'}$ if and only if they share the same weight in $\text{Wt}(Q)$ and the same image under $\tilde{\Phi}$.

Before introducing the main result, we present a technical lemma for any list \mathcal{L} as in (4.7). Write $\chi = \sum_i \chi_i \mathbf{e}_i \in \mathbb{Z}^{Q_0}$ as $\chi = \chi^+ - \chi^-$ where $\chi^\pm = \sum_i \chi_i^\pm \mathbf{e}_i \in \mathbb{N}^{Q_0}$ have disjoint supports $I_\chi^+ = \{i \in Q_0 : \chi_i > 0\}$ and $I_\chi^- = \{i \in Q_0 : \chi_i < 0\}$. In particular, $\chi \in \text{Wt}(Q)$ gives

$$n_\chi := \sum_{i \in I_\chi^+} \chi_i^+ = \sum_{i \in I_\chi^-} \chi_i^- \in \mathbb{N}.$$

For any spanning tree \mathcal{T} in Q , set $y_{\mathcal{T}} := \prod_{a \in \text{supp}(\mathcal{T})} y_a$. Recall that any path p in the quiver Q defines the monomial $y_p := \prod_{a \in \text{supp}(p)} y_a$.

Lemma 4.7. *Let \mathcal{T} be a spanning tree in Q and let $\chi \in \text{inc}(\mathbb{N}^{Q_1}) \setminus \{0\}$. There exists $\mathbf{m} \in \mathbb{N}^{Q_1}$ such that for any monomial $y^{\mathbf{v}} \in \mathbb{k}[y_a : a \in Q_1]$ of weight χ , we have*

$$(4.10) \quad (y_{\mathcal{T}})^{2n_{\chi}} y^{\mathbf{v}} \sim y^{\mathbf{m}} \prod_{\alpha=1}^{n_{\chi}} y_{\gamma_{\alpha}}$$

where $\gamma_1, \dots, \gamma_{n_{\chi}}$ are paths in Q , each with tail at 0 and head at r . Also, $y^{\mathbf{v}}$ divides $\prod_{\alpha=1}^{n_{\chi}} y_{\gamma_{\alpha}}$, and the resulting quotient $\tilde{\Phi}(\prod_{\alpha} y_{\gamma_{\alpha}}) / \tilde{\Phi}(y^{\mathbf{v}})$ depends only on \mathcal{T} and χ .

Proof. We begin by constructing $\mathbf{m} \in \mathbb{N}^{Q_1}$. The spanning tree \mathcal{T} supports a path q_i from $0 \in Q_0$ to each vertex $i \in Q_0$ and hence to each vertex in I_{χ}^{-} . We may therefore write

$$(4.11) \quad (y_{\mathcal{T}})^{n_{\chi}} = y^{\mathbf{m}_1} \prod_{i \in I_{\chi}^{-}} (y_{q_i})^{\chi_i^{-}}$$

where $\mathbf{m}_1 \in \mathbb{N}^{Q_1}$ depends only on \mathcal{T} and χ . The tree \mathcal{T} supports a path γ from 0 to r whose label is a torus-invariant section $s \in H^0(E_r)$. Since E_{r-1} is \mathcal{O}_X -regular with respect to E_1, \dots, E_{r-2} and each $\beta_i > 0$, Proposition 4.5 implies that the multiplication map

$$(4.12) \quad H^0(E_{r-1} \otimes E_1^{\delta\beta_1} \otimes \dots \otimes E_{r-2}^{\delta\beta_{r-2}} \otimes E_j^{-1}) \otimes_{\mathbb{k}} H^0(E_j) \rightarrow H^0(E_r)$$

is surjective. In particular, for each $j \leq r-2$ there exist sections of $E_r \otimes E_j^{-1}$ and E_j whose product is s . Since Q is a complete quiver of sections, there exists a pair of paths labelled by these sections, one from 0 to j denoted q_j'' , and the other from j to r denoted q_j' . Concatenating gives a path $q_j' q_j''$ from 0 to r that passes via j and, by Remark 4.6(2) through $r-1$, such that $y_{\gamma} \sim y_{q_j' q_j''} = y_{q_j'} y_{q_j''}$. Multiply by $y_{\mathcal{T}} / y_{\gamma} \in \mathbb{k}[y_a : a \in Q_1]$ to obtain $y_{\mathcal{T}} \sim y_{q_j'} y^{\mathbf{m}(j)}$ for some $\mathbf{m}(j) \in \mathbb{N}^{Q_1}$ that depends only on \mathcal{T} and j (and on the lift of s via (4.12), but we fix one such lift for \mathcal{T} and i). Applying this χ_j^{+} -times for each $j \in I_{\chi}^{+}$ and multiplying gives

$$(y_{\mathcal{T}})^{n_{\chi}} \sim y^{\mathbf{m}_2} \prod_{j \in I_{\chi}^{+}} (y_{q_j'})^{\chi_j^{+}}$$

for some $\mathbf{m}_2 \in \mathbb{N}^{Q_1}$ depends only on \mathcal{T} and χ . Multiply by (4.11) to see that

$$(4.13) \quad (y_{\mathcal{T}})^{2n_{\chi}} \sim y^{\mathbf{m}} \prod_{i \in I_{\chi}^{-}} (y_{q_i})^{\chi_i^{-}} \prod_{j \in I_{\chi}^{+}} (y_{q_j'})^{\chi_j^{+}}$$

where $\mathbf{m} := \mathbf{m}_1 + \mathbf{m}_2 \in \mathbb{N}^{Q_1}$ depends only on \mathcal{T} and χ .

To complete the proof, write $\mathbf{v} = \sum_{a \in Q_1} v_a \mathbf{e}_a \in \mathbb{N}^{Q_1}$ where $\text{inc}(\mathbf{v}) = \chi$. Since $\chi \neq 0$ there exists $i \in I_{\chi}^{-}$, so there exists $a_1 \in Q_1$ with $\mathbf{t}(a_1) = i$ such that $v_{a_1} > 0$. There are two cases. If $\chi_{\mathbf{h}(a_1)} < 0$ then $\mathbf{h}(a_1) \in I_{\chi}^{+}$, in which case we define $p_1 := a_1$ and repeat the above for $\mathbf{v}' := \mathbf{v} - \mathbf{e}_{a_1}$. Otherwise, $\chi_{\mathbf{h}(a_1)} \leq 0$ in which case there exists $a_2 \in Q_1$ with $\mathbf{t}(a_2) = \mathbf{h}(a_1)$ such that $v_{a_2} > 0$. Since Q is acyclic we can continue in this way, obtaining a path p_1 that traverses the arrows a_1, a_2, \dots and satisfies $\chi_{\mathbf{h}(p_1)} > 0$, that is, $\mathbf{h}(p_1) \in I_{\chi}^{+}$. As in the first case, we may repeat the above for $\mathbf{v}' := \mathbf{v} - \sum_{a \in \text{supp}(p_1)} \mathbf{e}_a$. In either case, the weight $\chi' := \text{inc}(\mathbf{v}')$ satisfies $n_{\chi'} = n_{\chi} - 1$, and we obtain by induction a set of paths $p_1, \dots, p_{n_{\chi}}$ satisfying $y^{\mathbf{v}} = \prod_{\alpha=1}^{n_{\chi}} y_{p_{\alpha}}$, where precisely χ_i^{-} of these paths have tail at $i \in I_{\chi}^{-}$ and χ_i^{+} have head at $i \in I_{\chi}^{+}$. Thus, for

$1 \leq \alpha \leq n_\chi$, there exists $i \in I_\chi^-, j \in I_\chi^+$ such that $\gamma_\alpha := q'_j p_\alpha q_i$ is a path in Q from 0 to r and

$$\prod_{\alpha=1}^{n_\chi} y_{\gamma_\alpha} = \prod_{i \in I_\chi^-} (y_{q_i})^{x_i^-} \prod_{\alpha=1}^{n_\chi} y_{p_\alpha} \prod_{i \in I_\chi^+} (y_{q'_i})^{x_i^+}.$$

Note that y^\vee divides $\prod_{\alpha=1}^{n_\chi} y_{\gamma_\alpha}$, and multiplying (4.13) by y^\vee establishes (4.10). The quotient $\tilde{\Phi}(\prod_{\alpha} y_{\gamma_\alpha}) / \tilde{\Phi}(y^\vee)$ equals $\tilde{\Phi}((y_\mathcal{T})^{2n_\chi}) / \tilde{\Phi}(y^\mathbf{m})$, so depends only on \mathcal{T} and χ as required. \square

Remark 4.8. Applying $\tilde{\Phi}(-)$ to (4.10) and dividing the resulting equality by $\tilde{\Phi}(y^\vee)$ shows in addition that the monomial $\tilde{\Phi}(y^\mathbf{m})$ divides $\tilde{\Phi}((y_\mathcal{T})^{2n_\chi})$.

We now come to our main algebraic result. The proof is rather involved, so we sketch the outline. The goal is to show that the ideals $I_\mathcal{L}, I_Q$ defined by some \mathcal{L} satisfy $I_Q = (I_\mathcal{L} : B_Q^\infty)$. This boils down to showing that for each $f \in I_Q$, there exists both $g \in I_\mathcal{L}$ and $N \in \mathbb{N}$ such that $(y_\mathcal{T})^N f \sim g$, where \sim is the equivalence relation from Remark 4.6(3). However, the generators of $I_\mathcal{L}$ from equation (4.2) are rather special polynomials of the form $\sum_{p \in \Gamma} c_p y_p$, where Γ is a finite set of paths with the same head and same tail, such that $\tilde{\Phi}(\sum_{p \in \Gamma} c_p y_p) \in I_X$. The goal, then, is to find an appropriate sets of paths $\{\Gamma_i\}$ for which an element $(y_\mathcal{T})^N f$ is equivalent under \sim to a polynomial combination of elements of the form $\sum_{p \in \Gamma_i} c_p y_p$ that satisfy $\tilde{\Phi}(\sum_{p \in \Gamma_i} c_p y_p) \in I_X$. STEP 1 below introduces sets of paths $\{\Gamma_i\}$ using Lemma 4.7, but at this stage we know only that $(y_\mathcal{T})^N f$ is equivalent under \sim to a combination of elements $\sum_{p \in \Gamma_i} c_p \prod_{\alpha} y_{p_\alpha}$. In STEP 2 we use multigraded regularity and the relation \sim to refine every such element so that, in STEP 3, we can pull out a sufficiently large common factor to leave generators of the form $\sum_{p \in \Gamma_i} c_p y_p$ as required. At each stage, we take care to control the appropriate element of I_X .

Theorem 4.9. *Let L_1, \dots, L_{r-2} be distinct, basepoint-free line bundles on X for which the corresponding reflexive sheaves $E_1 = \psi^{-1}(L_1), \dots, E_{r-2} = \psi^{-1}(L_{r-2})$ on \tilde{X} are basepoint-free line bundles. If the subsemigroup of $\text{Pic}(\tilde{X})$ generated by E_1, \dots, E_{r-2} contains an ample bundle, then there exists $L_{r-1}, L_r \in \text{Pic}(X)$ such that*

$$(4.14) \quad \varphi|_{\mathcal{L}} : X \longrightarrow \mathcal{M}_\vartheta(\text{mod-}A)$$

is an isomorphism for $\mathcal{L} = (\mathcal{O}_X, L_1, \dots, L_r)$ and $A = \text{End}_{\mathcal{O}_X}(\bigoplus_{0 \leq i \leq r} L_i)$.

Proof. Define the line bundles L_{r-1} and L_r as described at the start of this section to produce a collection \mathcal{L} of the form (4.7). Remark 4.6(1) shows that \mathcal{L} is very ample, so by Theorem 4.2 it suffices to prove that $I_Q = (I_\mathcal{L} : B_Q^\infty)$. One inclusion is straightforward: let $f \in (I_\mathcal{L} : B_Q^\infty)$. Since $I_\mathcal{L} \subseteq I_Q$ and hence $(I_\mathcal{L} : B_Q^\infty) \subseteq (I_Q : B_Q^\infty)$, we have that $(y_\mathcal{T})^N f \in I_Q$ for some spanning tree \mathcal{T} and $N \in \mathbb{N}$. Since I_Q is prime, we have either $f \in I_Q$ as required, or $B_Q \subseteq I_Q$. This latter conclusion would force the contradiction $\mathbb{V}(I_Q) \setminus \vartheta G = \emptyset$, so $(I_\mathcal{L} : B_Q^\infty) \subseteq I_Q$.

For the opposite inclusion, let $f \in I_Q$ be a homogeneous generator of weight $\chi \in \text{inc}(\mathbb{N}^{Q_1}) \setminus \{0\}$ and let \mathcal{T} be a spanning tree in Q . If we can show that $(y_\mathcal{T})^N f \in I_{\tilde{Q}} + I_\mathcal{L}$ for some $N \in \mathbb{N}$, then by increasing N if necessary and applying the equality $I_{\tilde{Q}} = (I_{\tilde{\mathcal{L}}} : B_{\tilde{Q}}^\infty)$ from (4.8), we deduce that $(y_\mathcal{T})^N f \in I_{\tilde{\mathcal{L}}} + I_\mathcal{L} = I_\mathcal{L}$ and hence $f \in (I_\mathcal{L} : B_Q^\infty)$ as required. In light of Remark 4.6(3), we need only find $g \in I_\mathcal{L}$ such that $(y_\mathcal{T})^N f \sim g$, and we achieve this for $N = 2n_\chi$.

We proceed in four steps:

STEP 1: Introduce a set of paths $\{\gamma_{\alpha,\beta}\}$ in Q such that

$$(4.15) \quad (y_{\mathcal{T}})^{2n_{\chi}} f \sim y^{\mathbf{m}} \left(\sum_{\beta} c_{\beta} \prod_{\alpha=1}^{n_{\chi}} y_{\gamma_{\alpha,\beta}} \right)$$

for some $\mathbf{m} \in \mathbb{N}^{Q^1}$ and $c_{\beta} \in \mathbb{k}$, where in addition we have $\tilde{\Phi}(\sum_{\beta} c_{\beta} \prod_{1 \leq \alpha \leq n_{\chi}} y_{\gamma_{\alpha,\beta}}) \in I_X$.

Decompose f as a sum of terms $f = \sum_{\beta} c_{\beta} y^{\mathbf{v}_{\beta}}$ for $c_{\beta} \in \mathbb{k}$ and $\mathbf{v}_{\beta} \in \mathbb{N}^{Q^1}$ satisfying $\chi = \text{inc}(\mathbf{v}_{\beta})$. Since $\chi \neq 0$ we apply Lemma 4.7 to each monomial $y^{\mathbf{v}_{\beta}}$ to obtain $(y_{\mathcal{T}})^{2n_{\chi}} y^{\mathbf{v}_{\beta}} \sim y^{\mathbf{m}} \prod_{\alpha=1}^{n_{\chi}} y_{\gamma_{\alpha,\beta}}$, where \mathbf{m} depends only on \mathcal{T} and χ (not on β) and where each $\gamma_{\alpha,\beta}$ is a path in Q with tail at 0 and head at r . This gives (4.15). Also, the quotient $x^{\mathbf{q}} := \tilde{\Phi}(\prod_{\alpha} y_{\gamma_{\alpha,\beta}}) / \tilde{\Phi}(y^{\mathbf{v}_{\beta}}) \in \mathbb{k}[x_1, \dots, x_d]$ depends only on \mathcal{T} and χ (not on β). Since $f \in I_Q$, we have $\tilde{\Phi}(f) \in I_X$ and hence we deduce that $\tilde{\Phi}(\sum_{\beta} c_{\beta} \prod_{\alpha=1}^{n_{\chi}} y_{\gamma_{\alpha,\beta}}) = x^{\mathbf{q}}(\sum_{\beta} c_{\beta} \tilde{\Phi}(y^{\mathbf{v}_{\beta}})) = x^{\mathbf{q}} \tilde{\Phi}(f) \in I_X$ as required.

STEP 2: Introduce a second set of paths $\{p_{i,j,k,\ell}\}$ in Q such that

$$\sum_{\beta} c_{\beta} \prod_{\alpha=1}^{n_{\chi}} y_{\gamma_{\alpha,\beta}} \sim \sum_{i,j,k} c_{i,j,k} \prod_{\ell=1}^{n_{\chi}} y_{p_{i,j,k,\ell}}$$

for some $c_{i,j,k} \in \mathbb{k}$, where for each i, j we have $\tilde{\Phi}(\sum_k c_{i,j,k} \prod_{1 \leq \ell \leq n_{\chi}} y_{p_{i,j,k,\ell}}) \in I_X$.

In light of Step 1, expand $\tilde{\Phi}(\sum_{\beta} c_{\beta} \prod_{\alpha=1}^{n_{\chi}} y_{\gamma_{\alpha,\beta}}) = \sum_{i,j} h_{i,j} g_i$ in terms of generators of I_X , where each $h_{i,j} \in \mathbb{k}[x_1, \dots, x_d]$ is a nonzero term. Since $\tilde{\Phi}$ is equivariant and $y_{\gamma_{\alpha,\beta}}$ has weight $\mathbf{e}_r - \mathbf{e}_0 \in \text{Wt}(Q)$, we may assume without loss of generality that each term in this expansion has degree $\widehat{\text{pic}}(n_{\chi}(\mathbf{e}_r - \mathbf{e}_0)) = E_r^{n_{\chi}}$. Thus, expanding each $g_i := g_{i,1} + \dots + g_{i,t_i}$ as a sum of terms for some $t_i \in \mathbb{N}$ gives $h_{i,j} g_{i,k} \in H^0(E_r^{n_{\chi}})$ for all i, j, k . Since E_{r-1} is \mathcal{O}_X -regular with respect to E_1, \dots, E_{r-2} and $E_r = E_{r-1}^2$, Proposition 4.5 implies that the multiplication map

$$H^0(E_r) \otimes_{\mathbb{k}} \dots \otimes_{\mathbb{k}} H^0(E_r) \rightarrow H^0(E_r^{n_{\chi}})$$

is surjective, so for each i, j, k there exists $c_{i,j,k} \in \mathbb{k}$ and torus-invariant sections $s_{i,j,k,\ell} \in H^0(E_r)$ for $1 \leq \ell \leq n_{\chi}$ such that $h_{i,j} g_{i,k} = c_{i,j,k} \prod_{\ell=1}^{n_{\chi}} s_{i,j,k,\ell}$. Since Q is a quiver of sections, there exists a path $p_{i,j,k,\ell}$ in Q from 0 to r whose label is the torus-invariant section $s_{i,j,k,\ell}$, that is, $\tilde{\Phi}(y_{p_{i,j,k,\ell}}) = s_{i,j,k,\ell}$. For fixed i, j , we therefore obtain

$$(4.16) \quad h_{i,j} g_{i,k} = c_{i,j,k} \tilde{\Phi} \left(\prod_{\ell=1}^{n_{\chi}} y_{p_{i,j,k,\ell}} \right).$$

Summing over $1 \leq k \leq t_i$ gives $h_{i,j} g_i = \tilde{\Phi}(\sum_k c_{i,j,k} \prod_{1 \leq \ell \leq n_{\chi}} y_{p_{i,j,k,\ell}})$, and by summing this new expression over all i, j we deduce that

$$(4.17) \quad \tilde{\Phi} \left(\sum_{\beta} c_{\beta} \prod_{\alpha=1}^{n_{\chi}} y_{\gamma_{\alpha,\beta}} \right) = \tilde{\Phi} \left(\sum_{i,j,k} c_{i,j,k} \prod_{\ell=1}^{n_{\chi}} y_{p_{i,j,k,\ell}} \right)$$

lies in I_X by Step 1. The main statement of Step 2 now follows from Remark 4.6(3) because these polynomials also share the same weight in $\text{Wt}(Q)$, namely $n_{\chi}(\mathbf{e}_r - \mathbf{e}_0)$.

STEP 3: Introduce a third set of paths $\{q_{i,j,k}\}$ in Q such that

$$(4.18) \quad \prod_{\ell=1}^{n_{\chi}} y_{p_{i,j,k,\ell}} \sim y^{\mathbf{m}'_{i,j}} y_{q_{i,j,k}}$$

for some $\mathbf{m}'_{i,j} \in \mathbb{N}^{Q_1}$, where for each i, j we have $\tilde{\Phi}(\sum_k c_{i,j,k} y_{q_{i,j,k}}) \in I_X$.

Fix i, j and simplify notation by suppressing dependence on i, j . To this end, write $c_k := c_{i,j,k}$, $y^{\mathbf{v}^k} := \prod_{1 \leq \ell \leq n_\chi} y_{p_{i,j,k,\ell}}$, $h = h_{i,j}$ and $g_k = g_{i,k}$, in which case (4.16) is simply $hg_k = c_k \tilde{\Phi}(y^{\mathbf{v}^k})$. The map $\tilde{\Phi}$ is equivariant and sends monomials to monomials, so $\frac{1}{c_k} hg_k \in H^0(E_r^{n_\chi})$ defines a torus-invariant section. Since $E = E_1^{\beta_1} \otimes \cdots \otimes E_{r-2}^{\beta_{r-2}}$ is \mathcal{O}_X -regular with respect to E_1, \dots, E_{r-2} and $E_r = E_{r-1}^2 = E^{2\delta}$, Proposition 4.5 implies that the multiplication map

$$H^0(E) \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} H^0(E) \rightarrow H^0(E_r^{n_\chi})$$

is surjective, so $\frac{1}{c_k} hg_k$ is equal to the product of $2\delta n_\chi$ torus-invariant sections of E . Since g_k is a term of a generator of I_X , its total degree is at most $\delta_0 \leq 2\delta$ by (4.6), so we may choose 2δ of these sections $s_{k,1}, \dots, s_{k,2\delta} \in H^0(E)$ such that g_k divides $\prod_{1 \leq \mu \leq 2\delta} s_{k,\mu} \in H^0(E_r)$. We now apply the above only for $k = 1$. Since Q is a quiver of sections, there exists a path q_1 in Q from 0 to r satisfying $\tilde{\Phi}(y_{q_1}) = \prod_{1 \leq \mu \leq 2\delta} s_{1,\mu}$, so the section $hg_1/c_1 \tilde{\Phi}(y_{q_1}) \in H^0(E_r^{n_\chi-1})$ is torus-invariant. Surjectivity of the multiplication map $H^0(E_r) \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} H^0(E_r) \rightarrow H^0(E_r^{n_\chi-1})$ determines $n_\chi - 1$ sections of E_r and hence paths $q'_1, \dots, q'_{n_\chi-1}$ in Q from 0 to r labelled by these sections such that $\tilde{\Phi}(y^{\mathbf{m}'}) = hg_1/c_1 \tilde{\Phi}(y_{q_1})$ for $y^{\mathbf{m}'} := \prod_{1 \leq \nu \leq n_\chi-1} y_{q'_\nu}$. In particular,

$$(4.19) \quad \tilde{\Phi}(y^{\mathbf{v}^1}) = \frac{hg_1}{c_1} = \tilde{\Phi}(y^{\mathbf{m}'} y_{q_1}).$$

Both monomials $y^{\mathbf{v}^1}$ and $y^{\mathbf{m}'} y_{q_1}$ have weight $n_\chi(\mathbf{e}_r - \mathbf{e}_0) \in \text{Wt}(Q)$, hence $y^{\mathbf{v}^1} \sim y^{\mathbf{m}'} y_{q_1}$. Now reintroduce the indices i, j , setting $\mathbf{m}'_{i,j} := \mathbf{m}'$ and $q_{i,j,1} := q_1$, to obtain (4.18) for $k = 1$.

For $k > 1$, we have $hg_k = c_k \tilde{\Phi}(y^{\mathbf{v}^k})$. For $1 \leq i \leq m$, the generator g_i of I_X is T -homogeneous, so $g_k := g_{i,k}$ and $g_1 := g_{i,1}$ have the same degree in \mathbb{Z}^ρ . Since g_1 divides $\tilde{\Phi}(y_{q_1}) \in H^0(E_r)$, it follows that the term $\tilde{\Phi}(y_{q_1}) g_k / g_1$ also has degree E_r . Divide by its coefficient $c_k / c_1 \in \mathbb{k}$ to obtain a torus-invariant section $\tilde{\Phi}(y_{q_1}) c_1 g_k / c_k g_1 \in H^0(E_r)$ which in turn determines a path q_k in Q with tail at 0 and head at r for which $\tilde{\Phi}(y_{q_k}) = \tilde{\Phi}(y_{q_1}) c_1 g_k / c_k g_1$. Then (4.19) gives

$$\tilde{\Phi}(y^{\mathbf{v}^k}) = hg_1 \cdot \frac{g_k}{c_k g_1} = c_1 \tilde{\Phi}(y^{\mathbf{m}'}) \tilde{\Phi}(y_{q_1}) \cdot \frac{g_k}{c_k g_1} = \tilde{\Phi}(y^{\mathbf{m}'} y_{q_k}).$$

It follows that the monomials $y^{\mathbf{v}^k}$ and $y^{\mathbf{m}'} y_{q_k}$ have weight $n_\chi(\mathbf{e}_r - \mathbf{e}_0)$, hence $y^{\mathbf{v}^k} \sim y^{\mathbf{m}'} y_{q_k}$, and we reintroduce the indices i, j , setting $q_{i,j,k} := q_k$, to obtain (4.18) for all k . Then

$$\tilde{\Phi}(y^{\mathbf{m}'_{i,j}}) \tilde{\Phi}\left(\sum_k c_{i,j,k} y_{q_{i,j,k}}\right) = \tilde{\Phi}\left(\sum_k c_{i,j,k} \prod_{\ell=1}^{n_\chi} y_{p_{i,j,k,\ell}}\right) \in I_X$$

holds for every i, j by combining (4.18) and Step 2. The ideal I_X does not contain the monomial $\tilde{\Phi}(y^{\mathbf{m}'_{i,j}})$ for any i, j , otherwise it would contain a variable of $\mathbb{k}[x_1, \dots, x_d]$ because I_X is prime, and we may assume this is not the case by redefining $\text{Cox}(X)$ from (2.1) using a polynomial ring in $d - 1$ variables. Thus, $\tilde{\Phi}(\sum_k c_{i,j,k} y_{q_{i,j,k}}) \in I_X$ for every i, j as required.

STEP 4: Establish that $(y_{\mathcal{T}})^{2n_\chi} f \in I_{\tilde{Q}} + I_{\mathcal{L}}$ as required by proving that

$$(4.20) \quad (y_{\mathcal{T}})^{2n_\chi} f \sim y^{\mathbf{m}} \left(\sum_{i,j} y^{\mathbf{m}'_{i,j}} \left(\sum_k c_{i,j,k} y_{q_{i,j,k}} \right) \right).$$

Relation (4.20) is immediate from Steps 1-3. For every i, j we also have $\sum_k c_{i,j,k} y_{q_{i,j,k}} \in I_{\mathcal{L}}$ by Step 3, so the right hand side of (4.20) also lies in $I_{\mathcal{L}}$. The definition of \sim given in Remark 4.6(3) then implies $(y_{\mathcal{T}})^{2n_X} f \in I_{\tilde{Q}} + I_{\mathcal{L}}$. This completes the proof of Theorem 4.9. \square

Example 4.10. Consider the Mori Dream Space $X = \text{Gr}(2, 4)$, where

$$\text{Cox}(X) = \frac{\mathbb{k}[x_1, \dots, x_6]}{(x_1x_6 - x_2x_5 + x_3x_4)}$$

and $\text{Pic}(X) \cong \mathbb{Z}$. Let L denote the very ample line bundle whose sections determine the Plücker embedding in the ambient toric variety $\tilde{X} = \mathbb{P}^5$. Consider Theorem 4.9 for $L_1 := L$. The generator of I_X has total degree 2, so by (4.6) we have $\delta = 1$. To construct the collection \mathcal{L} from (4.7), note that $E_1 = \mathcal{O}_{\mathbb{P}^5}(1)$ is normally generated and $\mathcal{O}_{\mathbb{P}^5}$ -regular with respect to E_1 . In this case, $E_{r-1} := E_1^\delta = E_1$ and we need only add $E_r := E^{2\delta} = \mathcal{O}_{\mathbb{P}^5}(2)$ to the collection to obtain $\tilde{\mathcal{L}} = (\mathcal{O}_{\mathbb{P}^5}, \mathcal{O}_{\mathbb{P}^5}(1), \mathcal{O}_{\mathbb{P}^5}(2))$ and hence $\mathcal{L} = (\mathcal{O}_X, L, L^2)$. The map $\varphi_{|\mathcal{L}|}: \text{Gr}(2, 4) \rightarrow |\mathcal{L}|$ is a closed immersion with image $\mathbb{V}(I_Q) //_{\vartheta} G$ by Proposition 3.5(i) and Theorem 3.7(iii). The quiver of sections Q is shown in Figure 2, where we list the arrows with tail at 0 as y_1, \dots, y_6




FIGURE 2. The quiver of sections Q for the collection \mathcal{L}

where $\text{div}(y_i) = x_i$ for $1 \leq i \leq 6$, and those with tail at 1 as y_7, \dots, y_{12} where $\text{div}(y_{i+6}) = x_i$ for $1 \leq i \leq 6$. The algorithm from Winn [15, Chapter 5] gives

$$I_{\mathcal{L}} = I_Q = \left(\begin{array}{l} y_1y_8 - y_2y_7, y_1y_9 - y_3y_7, y_1y_{10} - y_4y_7, y_1y_{11} - y_5y_7 \\ y_1y_{12} - y_6y_7, y_2y_9 - y_3y_8, y_2y_{10} - y_4y_8, y_2y_{11} - y_5y_8 \\ y_2y_{12} - y_6y_8, y_3y_{10} - y_4y_9, y_3y_{11} - y_5y_9, y_3y_{12} - y_6y_9 \\ y_4y_9 - y_2y_{11} + y_1y_{12}, y_4y_{11} - y_5y_{10}, y_4y_{12} - y_6y_{10}, y_5y_{12} - y_6y_{11} \\ y_9y_{10} - y_8y_{11} + y_7y_{12}, y_3y_{10} - y_2y_{11} + y_1y_{12}, y_3y_4 - y_2y_5 + y_1y_6 \end{array} \right),$$

so the natural inclusion map $\mathbb{V}(I_Q) //_{\vartheta} G \rightarrow \mathbb{V}(I_{\mathcal{L}}) //_{\vartheta} G$ from diagram (4.9) is an isomorphism. Thus, $\varphi_{|\mathcal{L}|}: \text{Gr}(2, 4) \rightarrow \mathcal{M}_{\vartheta}(\text{mod-}A)$ is an isomorphism for $A = \text{End}_{\mathcal{O}_X}(\bigoplus_{0 \leq i \leq 2} L_i)$.

Remark 4.11. We did not have to saturate $I_{\mathcal{L}}$ by B_Q in order to obtain I_Q in Example 4.10, but this is necessary for the choice $L_1 = L^2$. Indeed, the collection in this case is $\mathcal{L} = (\mathcal{O}_X, L^2, L^4)$, the ideals $I_{\mathcal{L}}$ and I_Q are computed explicitly by Winn [15, Section 6.4] and are shown to satisfy $I_Q = (I_{\mathcal{L}} : B_Q^\infty)$ even though $I_Q \neq I_{\mathcal{L}}$. The ideals have very many generators, so we do not reproduce them here. Instead we illustrate the saturation aspect of the construction with a pair of examples in the next section.

5. RECONSTRUCTING DEL PEZZO SURFACES FROM A TILTING BUNDLE

As an application of our main results we illustrate how to reconstruct del Pezzo surfaces directly from the bound quiver of sections of full, strongly exceptional collection of line bundles.

5.1. Tilting bundles on del Pezzo surfaces. Let X be a smooth projective variety over \mathbb{k} and write $\mathrm{coh}(X)$ for the category of coherent sheaves on X . For any vector bundle \mathcal{T} on X , let $A := \mathrm{End}_{\mathcal{O}_X}(\mathcal{T})$ denote its endomorphism algebra and $\mathrm{mod}\text{-}A$ the abelian category of finitely generated right A -modules. We say that \mathcal{T} is a *tilting bundle* on X if the functor

$$\mathrm{RHom}(\mathcal{T}, -) : D^b(\mathrm{coh}(X)) \longrightarrow D^b(\mathrm{mod}\text{-}A)$$

is an exact equivalence of bounded derived categories. If \mathcal{T} decomposes as a direct sum of line bundles $\mathcal{T} = \bigoplus_{0 \leq i \leq r} L_i$ (we need not assume that each L_i has rank one, but we choose to), then after reordering if necessary, the collection (L_0, L_1, \dots, L_r) is a full, strongly exceptional collection on X . That is, the line bundles in the collection generate $D^b(\mathrm{coh}(X))$ as a triangulated category and they satisfy appropriate Ext-vanishing conditions, namely, that $\mathrm{Hom}(L_j, L_i) = 0$ for $j > i$ and that $\mathrm{Ext}^k(L_i, L_j) = 0$ for $k > 0$ and all $0 \leq i, j \leq r$.

For $0 \leq k \leq 8$, let X_k denote the del Pezzo surface obtained as the blow-up of $\mathbb{P}_{\mathbb{k}}^2$ at k points in general position. The Picard group \mathbb{Z}^{k+1} has a basis given by H , the pullback to X_d of the hyperplane class on $\mathbb{P}_{\mathbb{k}}^2$, together with the k exceptional curves R_1, \dots, R_k . Consider the sequence of basepoint-free line bundles

$$(5.1) \quad \mathcal{L}_k := (\mathcal{O}_{X_k}, H, 2H - R_1, \dots, 2H - R_k, 2H)$$

on X_k , and write $L_0 = \mathcal{O}_{X_k}$, $L_1 = H$, $L_{i+1} = 2H - R_i$ for $1 \leq i \leq k$, and $L_{k+2} = 2H$.

Lemma 5.1. *The collection \mathcal{L}_k is very ample for $0 \leq k \leq 8$.*

Proof. For $k = 4$, the collection \mathcal{L}_4 is precisely the collection (3.8), and $\varphi_{|\mathcal{L}_4|}$ was shown to be a closed immersion in Example 3.10. The proof in that example is valid for any $0 \leq k \leq 8$. \square

Remark 5.2. In the terminology of Bergman–Proudfoot [3], Lemma 5.1 asserts that ϑ is *great*.

The following result is well known to experts, but for convenience we guide the reader towards a proof using the technology of toric systems introduced by Hille–Perling [10]. A *toric system* is a (cyclically ordered) list of Cartier divisors whose sum is the anticanonical divisor, such that the intersection product of any two adjacent divisors equals 1 and the intersection product of any distinct and non-adjacent divisors equals 0.

Proposition 5.3. *The line bundles (5.1) on X_k form a full, strongly exceptional collection, that is, the vector bundle $\mathcal{T}_k := \bigoplus_{0 \leq i \leq k+2} L_i$ is tilting.*

Proof. Beginning with the unique toric system H, H, H on $\mathbb{P}_{\mathbb{k}}^2$, construct a toric system on each X_k as follows: choose $H, H - R_1, R_1, H - R_1$ on X_1 , repeat for $k \geq 2$, introducing R_k in the second-last position while subtracting R_k from each neighbouring divisor to obtain the system

$$H, H - R_1, R_1 - R_2, R_2 - R_3, \dots, R_{k-1} - R_k, R_k, H - \sum_{1 \leq i \leq k} R_i$$

on X_k . List these divisors from left to right as D_1, \dots, D_{k+3} . Observe that for $1 \leq i \leq k+2$ we have $L_i = \mathcal{O}(D_1 + \dots + D_i)$, and $-K_{X_k} = \mathcal{O}(D_1 + \dots + D_{k+3})$, and Hille–Perling [10, Theorem 5.7] establishes that $(L_0, L_1, \dots, L_{k+2})$ is full, strongly exceptional sequence as required. \square

Thus far then, the variety X_k is smooth, the vector bundle \mathcal{T}_k is tilting, and the morphism $\varphi_{|\mathcal{L}_k|} : X_k \rightarrow |\mathcal{L}_k|$ is a closed immersion. A result of Bergman–Proudfoot [3, Theorem 2.4] now shows that $\varphi_{|\mathcal{L}_k|}$ identifies X_k with a connected component of the appropriate moduli space:

Corollary 5.4. *For $0 \leq k \leq 8$ and for the algebra $A_k := \mathbb{k}Q/J_{\mathcal{L}_k}$, the morphism $\varphi|_{\mathcal{A}}$ identifies X_k with a connected component of the moduli space $\mathcal{M}_\vartheta(\text{mod-}A_k)$.*

We now apply our main results to refine the statement of Corollary 5.4. Let (Q_k, J_k) denote the bound quiver of sections of the collection \mathcal{L}_k on X_k . For $k \leq 3$, the variety X_k is a toric variety, in which case $\mathcal{L}_k = \widetilde{\mathcal{L}}_k$ and the method of Craw–Smith [6] shows that the morphism $\varphi|_{\mathcal{L}_k}: X_k \rightarrow \mathcal{M}_\vartheta(\text{mod-}A_k)$ is an isomorphism in each case. The Mori Dream Space X_k is not toric for $k > 3$ and we now investigate a pair of examples.

5.2. The four-point blow-up. On the del Pezzo X_4 , the collection

$$\mathcal{L}_4 := (\mathcal{O}_{X_4}, H, 2H - R_1, 2H - R_2, 2H - R_3, 2H - R_4, 2H)$$

from (5.1) is precisely the collection listed in (3.8), and the quiver of sections Q is shown in Figure 1. In this case, the irrelevant ideal in \mathbb{A}^{Q_1} is given by the intersection of monomial ideals

$$B_Q = (y_1, \dots, y_6) \cap (y_7, y_8, y_9) \cap (y_{10}, y_{11}, y_{12}) \cap (y_{13}, y_{14}, y_{15}) \cap (y_{16}, y_{17}, y_{18}) \cap (y_{19}, \dots, y_{22}),$$

and the ideal I_Q is given in Example 3.10. Winn [15, Chapter 5] computes

$$I_{\mathcal{L}_4} = \left(\begin{array}{l} y_{15}y_{21} - y_{18}y_{22}, y_{12}y_{20} - y_{17}y_{22}, y_{11}y_{20} - y_{14}y_{21}, y_9y_{19} - y_{16}y_{22} \\ y_8y_{19} - y_{13}y_{21}, y_7y_{19} - y_{10}y_{20}, y_6y_{17} - y_5y_{18}, y_6y_{16} - y_3y_{18} \\ y_5y_{16} - y_3y_{17}, y_6y_{14} - y_4y_{15}, y_6y_{13} - y_2y_{15}, y_4y_{13} - y_2y_{14}, y_5y_{11} - y_4y_{12} \\ y_5y_{10} - y_1y_{12}, y_4y_{10} - y_1y_{11}, y_3y_8 - y_2y_9, y_3y_7 - y_1y_9, y_2y_7 - y_1y_8 \\ y_3y_{14}y_{21} - y_4y_{16}y_{22}, y_5y_{13}y_{21} - y_2y_{17}y_{22}, y_6y_{10}y_{20} - y_1y_{18}y_{22} \\ y_{16} - y_{17} + y_{18}, y_{13} - y_{14} + y_{15}, y_{10} - y_{11} + y_{12}, y_7 - y_8 + y_9, y_3 - y_5 + y_6 \\ y_2 - y_4 + y_6, y_1 - y_4 + y_5, y_{15}y_{21} - y_{18}y_{22}, y_{12}y_{20} - y_{17}y_{22}, y_{11}y_{20} - y_{14}y_{21} \\ y_9y_{19} - y_{17}y_{22} + y_{18}y_{22}, y_8y_{19} - y_{14}y_{21} + y_{18}y_{22}, y_6y_{17} - y_5y_{18} \\ y_6y_{14} - y_4y_{15}, y_5y_{11} - y_4y_{12}, y_5y_8 - y_6y_8 - y_4y_9 + y_6y_9 \end{array} \right),$$

and a Macaulay2 [8] computation gives $I_Q = (I_{\mathcal{L}_4} : B_Q^\infty)$ in this case. Since \mathcal{L}_4 is very ample, Theorem 4.2 implies that the closed immersion $\varphi|_{\mathcal{L}_4}: X_4 \rightarrow \mathcal{M}_\vartheta(\text{mod-}A_4)$ is an isomorphism.

5.3. The five-point blow-up. To obtain X_5 we blow-up $\mathbb{P}_{\mathbb{k}}^2$ at the torus-invariant points in addition to $(1, 1, 1)$ and $(1, 2, 3)$, in which case $\text{Cox}(X_5)$ is the quotient of $\mathbb{k}[x_1, \dots, x_{16}]$ by

$$I_{X_5} = \left(\begin{array}{l} x_5x_{16} + x_6x_{13} - 3x_8x_{10}, x_4x_{16} + 2x_6x_{14} + x_7x_{12}, x_4x_{16} + x_6x_{14} + x_9x_{10} \\ x_3x_{16} + x_6x_{15} + x_8x_{12}, x_3x_{16} + 2x_6x_{15} + x_9x_{11}, x_2x_{16} + x_7x_{15} - 2x_8x_{14} \\ x_2x_{16} + 3x_7x_{15} + 2x_9x_{13}, x_1x_{16} + 2x_{10}x_{15} - x_{11}x_{14}, x_1x_{16} + 3x_{10}x_{15} + x_{12}x_{13} \\ x_2x_6 - x_3x_7 + x_4x_8, 2x_2x_6 - 3x_3x_7 - x_5x_9, x_1x_6 - x_3x_{10} + x_4x_{11} \\ x_1x_6 - 3x_3x_{10} - x_5x_{12}, x_1x_7 - x_2x_{10} + x_4x_{13}, x_1x_7 - 2x_2x_{10} + x_5x_{14} \\ x_1x_8 - x_2x_{11} + x_3x_{13}, -2x_1x_8 + x_2x_{11} - x_5x_{15}, -x_1x_9 + 2x_2x_{12} + 3x_3x_{14} \\ -2x_1x_9 + x_2x_{12} + 3x_4x_{15}, x_6x_{13} - x_7x_{11} + x_8x_{10} \end{array} \right),$$

see Winn [15, Chapter 2]. On X_5 , the full strongly exceptional collection (5.1) is

$$\mathcal{L}_5 := (\mathcal{O}_{X_5}, H, 2H - R_1, 2H - R_2, 2H - R_3, 2H - R_4, 2H - R_5, 2H)$$

and the quiver of sections Q is shown in Figure 3 (in fact we omit one arrow labelled $x_1x_2x_4x_5x_{16}$ with tail at 0 and head at 4 to prevent the figure from becoming illegible). Arrows with tail at 0 and head at 1 are listed a_1, \dots, a_{10} from the top of Figure 3 to the bottom; list those with




FIGURE 3. Almost the quiver of sections for \mathcal{L}_5 (one arrow omitted from 0 to 4)

tail at 1 as a_{11}, \dots, a_{30} from top to bottom; list those with head at 7 as a_{31}, \dots, a_{35} from top to bottom; and list those with tail at 0 and head at $i \geq 2$ as a_{36}, \dots, a_{40} from top to bottom, where the arrow omitted from the figure is a_{38} . List the coordinates of $\mathbb{A}_{\mathbb{k}}^{Q_1}$ as y_1, \dots, y_{40} , and compute the kernel of (3.4) to obtain the ideal

$$I_Q = \left(\begin{array}{l} y_7 + 2y_9 - y_{10}, y_6 - 2y_8 + y_{10}, y_5 + y_8 - y_9, y_4 + y_9 - 2y_{10}, y_3 - y_8 + y_1 \\ y_2 + 2y_8 - y_9, y_1 + 3y_8 - y_9 - y_{10}, y_{28} + 2y_{29} - y_{30}, y_{27} + y_{29} - 2y_{30} \\ y_{24} - 2y_{25} + y_{26}, y_{23} - y_{25} + y_{26}, y_{20} + y_{21} - y_{22} \\ y_{19} + 2y_{21} - y_{22}, 2y_{16} + y_{17} + y_{18}, 2y_{15} + 3y_{17} + y_{18} \\ y_{12} + 2y_{13} + y_{14}, y_{11} + 3y_{13} + y_{14}, 10y_{29} - y_9y_{30}, y_{14}y_{31} + y_9 - 2y_{10} \\ 2y_8y_{29} + 2y_8y_{30} - 3y_9y_{30} - y_{40}, y_{10}y_{25} - y_8y_{26}, y_{13}y_{31} - y_8 + y_{10} \\ 2y_9y_{25} + 2y_8y_{26} - 3y_9y_{26} - y_{39}, 2y_{10}y_{21} + 2y_8y_{22} - 3y_{10}y_{22} - y_{38} \\ y_9y_{21} - y_8y_{22}, 3y_{10}y_{17} - 2y_8y_{18} + 3y_{10}y_{18} - 2y_{37}, y_{21}y_{26}y_{29} - y_{22}y_{25}y_{30} \\ 3y_9y_{17} + 2y_8y_{18} - y_{37}, 6y_{10}y_{13} - 2y_8y_{14} + 3y_{10}y_{14} - y_{36} \\ 3y_9y_{13} + y_8y_{14} - y_{36}, y_{30}y_{35} - y_{10}, y_{29}y_{35} - y_9, y_{26}y_{34} - y_{10} \\ y_{25}y_{34} - y_8, y_{22}y_{33} - y_9, y_{21}y_{33} - y_8, y_{18}y_{32} + 2y_9 - y_{10}, y_{17}y_{32} - 2y_8 + y_{10} \end{array} \right).$$

Details of the computation are given by Winn [15, Chapter 6], where it is also shown that

$$I_{\mathcal{L}_5} = \left(\begin{array}{l} y_7 + 2y_9 - y_{10}, y_6 - 2y_8 + y_{10}, y_5 + y_8 - y_9, y_4 + y_9 - 2y_{10}, y_3 - y_8 + y_{10} \\ y_2 + 2y_8 - y_9, y_1 + 3y_8 - y_9 - y_{10}, y_{28} + 2y_{29} - y_{30}, y_{27} + y_{29} - 2y_{30} \\ y_{24} - 2y_{25} + y_{26}, y_{23} - y_{25} + y_{26}, y_{20} + y_{21} - y_{22}, y_{19} + 2y_{21} - y_{22} \\ 2y_{16} + y_{17} + y_{18}, 2y_{15} + 3y_{17} + y_{18}, y_{12} + 2y_{13} + y_{14}, y_{11} + 3y_{13} + y_{14} \\ y_{10}y_{29} - y_9y_{30}, 2y_8y_{29} + 2y_8y_{30} - 3y_9y_{30} - y_{40}, y_{10}y_{25} - y_8y_{26} \\ 2y_9y_{25} + 2y_8y_{26} - 3y_9y_{26} - y_{39}, 2y_{10}y_{21} + 2y_8y_{22} - 3y_{10}y_{22} - y_{38} \\ y_9y_{21} - y_8y_{22}, 3y_{10}y_{17} - 2y_8y_{18} + 3y_{10}y_{18} - 2y_{37}, 3y_9y_{17} + 2y_8y_{18} - y_{37} \\ 6y_{10}y_{13} - 2y_8y_{14} + 3y_{10}y_{14} - y_{36}, 3y_9y_{13} + y_8y_{14} - y_{36}, y_{26}y_{34} - y_{30}y_{35} \\ y_{22}y_{33} - y_{29}y_{35}, y_{21}y_{33} - y_{25}y_{34}, y_{18}y_{32} + 2y_{29}y_{35} - y_{30}y_{35} \\ y_{17}y_{32} - 2y_{25}y_{34} + y_{30}y_{35}, y_{14}y_{31} + y_{29}y_{35} - 2y_{30}y_{35} \\ y_{13}y_{31} - y_{25}y_{34} + y_{30}y_{35} \end{array} \right).$$

A computation using Macaulay2 gives that $I_Q = (I_{\mathcal{L}_5} : B_Q^\infty)$ in this case. Since \mathcal{L}_5 is very ample, Theorem 4.2 implies that the morphism $\varphi_{|\mathcal{L}_5|} : X_5 \rightarrow \mathcal{M}_\vartheta(\text{mod-}A_5)$ is an isomorphism.

REFERENCES

- [1] Victor Batyrev and Oleg Popov. The Cox ring of a del Pezzo surface. In *Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002)*, volume 226 of *Progr. Math.*, pages 85–103. Birkhäuser Boston, Boston, MA, 2004.
- [2] Florian Berchtold and Jürgen Hausen. Cox rings and combinatorics. *Trans. Amer. Math. Soc.*, 359(3):1205–1252 (electronic), 2007.
- [3] Aaron Bergman and Nicholas Proudfoot. Moduli spaces for Bondal quivers. *Pacific J. Math.*, 237(2):201–221, 2008.
- [4] David A. Cox, John B. Little, and Henry K. Schenck. *Toric varieties*, volume 124 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.
- [5] Alastair Craw. Quiver flag varieties and multigraded linear series. *Duke. Math. J.*, 156(3):469–500, 2011.
- [6] Alastair Craw and Gregory G. Smith. Projective toric varieties as fine moduli spaces of quiver representations. *Amer. J. Math.*, 130(6):1509–1534, 2008.
- [7] E. Javier Elizondo, Kazuhiko Kurano, and Kei-ichi Watanabe. The total coordinate ring of a normal projective variety. *J. Algebra*, 276(2):625–637, 2004.
- [8] Daniel Grayson and Michael Stillman. Macaulay 2, a software system for research in algebraic geometry. Available from <http://www.math.uiuc.edu/Macaulay2/>.
- [9] Milena Hering, Hal Schenck, and Gregory G. Smith. Syzygies, multigraded regularity and toric varieties. *Compos. Math.*, 142(6):1499–1506, 2006.
- [10] Lutz Hille and Markus Perling. Exceptional sequences of invertible sheaves on rational surfaces. *Compos. Math.*, 147(4):1230–1280, 2011.
- [11] Yi Hu and Sean Keel. Mori dream spaces and GIT. *Michigan Math. J.*, 48:331–348, 2000. Dedicated to William Fulton on the occasion of his 60th birthday.
- [12] Alastair King. Moduli of representations of finite-dimensional algebras. *Quart. J. Math. Oxford Ser. (2)*, 45(180):515–530, 1994.
- [13] Antonio Laface and Mauricio Velasco. A survey on Cox rings. *Geom. Dedicata*, 139:269–287, 2009.
- [14] Diane Maclagan and Gregory G. Smith. Multigraded Castelnuovo-Mumford regularity. *J. Reine Angew. Math.*, 571:179–212, 2004.
- [15] Dorothy Winn. *Mori Dream Spaces as fine moduli of quiver representations*, 2011. University of Glasgow PhD thesis, 93pp, June 2011.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GLASGOW, GLASGOW, G12 8QW, UNITED KINGDOM
E-mail address: Alastair.Craw@glasgow.ac.uk, Dorothy.Winn@glasgow.ac.uk